

## Research Article



## Effect of Fraction 3 of *portulaca oleracea* on Haematological and Plasma Biochemical Parameters in Male Albino Rats

Oyededeji K.O.<sup>1</sup>, Adegoke A.O.<sup>2</sup>, Oladosu I.A.<sup>3</sup>

<sup>1</sup>Department of Physiology, College of Medicine and Health Sciences, Afe Babalola University, Ado-Ekiti, Nigeria.

<sup>2</sup>Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Ibadan, Ibadan, Nigeria.

<sup>3</sup>Department of Chemistry, Faculty of Science, University of Ibadan, Ibadan, Nigeria.

\*Corresponding author's E-mail: [sinaoyededeji@yahoo.com](mailto:sinaoyededeji@yahoo.com)

Accepted on: 05-06-2013; Finalized on: 30-06-2013.

### ABSTRACT

The effect of oral administration of chromatographic fraction 3 of *Portulaca oleracea* at doses of 1 mg/kg BW, 2 mg/kg BW and 3 mg/kg BW on haematological and plasma biochemical parameters of albino rats were investigated. The fraction was administered on daily basis for 30 days and blood samples were collected for analyses. Treatment of rats with 3 mg/kg BW of fraction 3 caused significant ( $p < 0.05$ ) decrease in the PCV and RBC values relative to their respective controls, while treatment of rats with 3 mg/kg BW of fraction 3 caused significant ( $p < 0.05$ ) increases in TWBC and platelet counts relative to their respective controls. Treatment of rats with 1 mg/kg BW, 2 mg/kg BW and 3 mg/kg of fraction 3 resulted in significant ( $p < 0.05$ ) increases in the albumin level as well as AST and ALT activities relative to their respective controls. These findings on haematological and plasma biochemical parameters suggest that the possible changes in blood chemistry of the treated rats were due to chromatographic fraction 3 of *Portulaca oleracea*.

**Keywords:** Albino rats, Albumin, Chromatographic fraction 3, *Portulaca oleracea*, Red blood cell.

### INTRODUCTION

*Portulaca oleracea* belongs to the family of *Portulacaceae*. It is commonly called Purslane in English language, "Babbajibi" in Hausa language and "Esan omode" or "Papasan" in Yoruba language. It is a fleshy annual herb, much-branched and attaining 30 cm long.<sup>1</sup>

It is used medicinally in Ghana for heart-palpitations.<sup>2</sup> The plant is used as a diuretic in Nigeria.<sup>3</sup> A tisane of the plant is drunk in Trinidad as a vermifuge.<sup>4</sup> At some areas near Benin City (Nigeria), the plant, along with other ingredients is taken as an aid to the development of foetus.<sup>5</sup>

It has been reported that chromatographic fractions of *Portulaca oleracea* have antifertility effects in male albino rats.<sup>6</sup> It has been reported that crude extracts of *Portulaca oleracea* have deleterious effect on the reproductive functions in male albino rats.<sup>7</sup> Isolated lupone constituent of *Portulaca oleracea* has also been reported to have deleterious effect on reproductive functions in male albino rats.<sup>8</sup>

This study aims at investigating the effect of chromatographic fraction 3 of *Portulaca oleracea* on the haematological and plasma biochemical parameters in male albino rats.

### MATERIALS AND METHODS

#### Experimental Animals

Adult male albino rats weighing between 160 g and 180 g bred in the Animal House of Physiology Department, LAUTECH, Ogbomoso were used. They were housed

under standard laboratory conditions with a 12 hours daylight cycle and had free access to feed and water; they were acclimatized to laboratory conditions for two weeks before the commencement of the experiments. All experiments were carried out in compliance with the recommendations of Helsinki's declaration on guiding principles on care and use of animals.

#### Plant Material

Fresh specimens of *Portulaca oleracea* were collected from the Botanical Garden of the Forestry Research Institute of Nigeria, Jericho, Ibadan, and was authenticated in the above named institute where a voucher specimen (No FHI 108334) was deposited.

#### Extraction and Fractionation of *Portulaca oleracea*

About 3.2 kg of air-dried specimen of *Portulaca oleracea* was cold-extracted in methanol for 72 hours. The mixture was filtered using a wire-gauze and a sieve with tiny pores (0.25 mm) and concentrated at room temperature by exposing the extract for six days. The resulting solution was then placed in the oven at a reduced temperature (50°C).

The methanolic extract was then preabsorbed with silica gel and placed in the oven at a reduced temperature (50°C) overnight and then subjected to open column chromatography on silica gel (F<sub>254</sub>, 50-200 mesh, E. Merck) for fractionation. The solvents (mobile phases) were hexane (non-polar), ethylacetate (partially polar) and methanol (polar). The gradients of the mobile phases involved hexane with an increasing percentage of ethylacetate (hexane/ethylacetate mixture) and then



ethylacetate with an increasing percentage of methanol (ethylacetate/methanol mixture).

### Thin Layer Chromatography (TLC)

The 21 fractions were spotted on precoated plates of silica gel GF<sub>254</sub> (20 x 20, 0.5 mm thick; E. Merck) using capillary tubes. The spotted TLC plates were developed in a tank that contained a mixture of ethylacetate/methanol (9:1) as the mobile phases.

The TLC plates were then examined under the ultraviolet (UV) light at a wavelength of 365 nm and the well-defined spots of the components were then revealed by the UV light. Fractions with similar relative fronts or retention or retardation factors ( $R_f$  value) were then pooled or bulked together, this then reduced the number of fractions to five (fractions 1, 2, 3, 4, 5)

$$R_f = \frac{\text{Distance compound has moved from origin}}{\text{Distance of solvent front from origin}}$$

Fraction 3 was then subjected to bioassay, vis-à-vis, its effect on haematological and plasma biochemical parameters in male albino rats were evaluated.

The dosages of the fraction administered in this study were extrapolated from that reported by.<sup>9</sup>

### Experimental Design

Twenty animals were randomly divided into four groups with each group consisting of five rats. The four groups were subjected to the following oral daily treatments for 30 days:

Group I rats received 1 mg/kg BW of fraction 3

Group II rats received 2 mg/kg BW of fraction 3

Group III rats receive 3 mg/kg BW of fraction 3

Group IV rats received 0.5 ml of distilled water as the control group.

### Collection of Blood Samples

Blood samples were collected through the medial cantus into EDTA bottles for haematological and plasma biochemical studies. Before assays, the blood samples were centrifuge for 5 minutes using a bench-top centrifuge (Centromix) and the supernant plasma was then used for the determinations of the biochemical parameters.

### Determination of Haematological Parameters

The red blood cells (RBC) and white blood cells (WBC) counts were determined by the improved Neubauer haemocytometer method. The haemoglobin (Hb) concentration was determined according to<sup>10</sup>, using the cyanomethaemoglobin method. The packed cell volume (PCV) was determined by the micro-haematocrit method according to.<sup>11</sup> Schilling method of differential leucocyte count was used to determine the distribution of the various white blood cells.<sup>12</sup> Mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH) and mean

corpuscular haemoglobin concentration (MCHC) were computed according to.<sup>10</sup>

### Determination of Plasma Biochemical Parameters

The total protein concentration was determined using the Biuret method<sup>13</sup> and the albumin concentration by the method of.<sup>14</sup> The globulin concentration was calculated by subtracting the albumin concentration from the total protein concentration. Activities of plasma alanine transaminase (ALT) and aspartate transaminase (AST) were determined according to the method of.<sup>15</sup> All the above biochemical parameters were determined in the plasma using the Randox kits.

### Statistical Analysis

The mean and standard error of mean (S.E.M.) were calculated for all values. Comparison between the control and experimental groups was done using one-way analysis of variance (ANOVA) with Duncan's Multiple Range Test. Differences were considered statistically significant at  $p < 0.05$ .

## RESULTS

The effects of various doses of fraction on haematological and plasma biochemical parameters of albino rats after treatment of rats for 30 days are shown respectively in Tables 1 and 2.

Treatment of rats with 3 mg/kg BW of fraction 3 caused significant ( $p < 0.05$ ) decrease in the PCV and RBC values relative to their respective controls, while treatment of rats with 3 mg/kg BW of fraction 3 caused significant ( $p < 0.05$ ) increases in TWBC and platelet counts relative to their respective controls. Treatment of rats with 2 mg/kg BW of fraction 3 caused significant ( $p < 0.05$ ) increase and decrease respectively in neutrophil and lymphocyte counts relative to their respective controls. Treatment of rats with 3 mg/kg BW of fraction 3 caused significant ( $p < 0.05$ ) decrease in monocyte count relative to the control. Treatment of rats with 1 mg/kg BW of fraction 3 caused significant ( $p < 0.05$ ) increases in MCHC and MCH values relative to their respective controls. Treatment of rats with 1 mg/kg BW, 2 mg/kg BW and 3 mg/kg BW of fraction 3 caused non-significant ( $p > 0.05$ ) changes in MCV, Hb and eosinophil values relative to their respective control.

Treatment of rats with 1 mg/kg BW, 2 mg/kg BW and 3 mg/kg BW of fraction 3 resulted in significant ( $p < 0.05$ ) increases in the albumin level as well as AST and ALT activities relative to their respective controls. Treatment of rats with 2 mg/kg BW of fraction 3 caused significant ( $p < 0.05$ ) increases in total protein and globulin levels relative to their respective controls.

## DISCUSSION

Fraction 3 caused significant decrease in PCV values, this could indicate an induction of anaemia.<sup>16</sup> Similar results were reported by<sup>17</sup> in fractions 3 and 4 of *Cnestis ferruginea* treated rats.

**Table 1:** Effect of Varying Doses of Fraction 3 on Haematological Parameters after Treatment of Rats for 30 Days

Parameters	Control	1 mg/kg	2 mg/kg	3 mg/kg
PCV (%)	41.80 ± 2.32	43.00 ± 1.47	43.00 ± 0.71	37.00 ± 1.08*
Hb (g/dl)	13.10 ± 0.83	14.30 ± 0.48	13.80 ± 0.32	12.00 ± 0.22
RBC (x10 <sup>6</sup> /μl)	7.04 ± 0.39	7.10 ± 0.13	7.24 ± 0.05	6.27 ± 0.18*
MCV (FL)	59.20 ± 0.34	60.50 ± 1.01	59.40 ± 0.78	59.00 ± 0.31
MCHC (g/dl)	31.30 ± 0.27	33.30 ± 0.03*	32.20 ± 0.26	32.40 ± 0.40
MCH (pg)	18.60 ± 0.26	20.10 ± 0.32*	19.10 ± 0.40	19.10 ± 0.28
TWBC (x10 <sup>3</sup> /μL)	8.34 ± 0.62	7.64 ± 0.74	9.00 ± 1.30	11.00 ± 0.13*
Platelets (10 <sup>5</sup> /μL)	1.20 ± 0.13	1.10 ± 0.03	1.60 ± 0.24*	2.00 ± 0.03*
Neutrophils (%)	32.80 ± 1.93	34.80 ± 2.50	50.80 ± 3.68*	35.80 ± 2.90
Lymphocytes (%)	64.00 ± 1.41	62.30 ± 2.72	45.80 ± 2.87*	62.80 ± 3.28
Eosinophils (%)	1.75 ± 0.48	2.00 ± 0.41	2.00 ± 0.41	1.25 ± 0.63
Monocytes (%)	2.00 ± 0.41	1.00 ± 0.41	1.25 ± 0.63	0.25 ± 0.25*

(n = 5, \*p&lt;0.05)

**Table 2:** Effect of Varying Doses of Fraction 3 on Serum Biochemical Parameters after Treatment of Rats for 30 Days

Parameters	Control	1 mg/kg	2 mg/kg	3 mg/kg
Total Protein (gm%)	4.90 ± 0.18	5.18 ± 0.14	6.70 ± 0.07*	4.93 ± 0.12
Albumin (gm%)	1.28 ± 0.10	1.73 ± 0.15*	2.23 ± 0.05*	1.68 ± 0.09*
Globulin (gm%)	3.63 ± 0.10	3.43 ± 0.09	4.48 ± 0.10*	3.93 ± 0.35
AST (μ/L)	19.30 ± 2.32	25.00 ± 1.63*	35.30 ± 1.44*	32.00 ± 1.08*
ALT (μ/L)	14.80 ± 1.84	22.30 ± 2.06*	29.00 ± 0.82*	29.80 ± 1.03*

(n = 5, \*p&lt;0.05)

Fraction 3 caused significant reductions in RBC counts, which might also indicate the induction of anaemia. Similar results were reported by<sup>18</sup> in *A. Cordifolia*, *P. amarus*, *P. muellerianus* and *S. virosa* extracts treated rats. Fraction 3 caused significant increase in MCH values which probably indicates the induction of macrocytic anaemia, since increased MCV and MCH values are known to be indicative of macrocytic anaemia.<sup>19</sup> Similar result was reported by<sup>20</sup> in *Azadirachta indica* extract treated chicks. The significant increase in the MCHC values caused by fraction 3 could indicate induction of hereditary spherocytosis, since MCHC values are known to be elevated in hereditary spherocytosis. Contrary results were reported by<sup>21</sup> in Garlic and Ginger supplements fed broilers.

Fraction 3 caused significant increase in TWBC value, this probably indicates an enhancement in the ability of the body to defend against invading organisms.<sup>22</sup> Similar result was reported by<sup>23</sup> in *viscum album* extract treated rats. Fraction 3 caused significant decrease in monocyte values which indicates that the phagocytic function of the body could be compromised.<sup>24</sup> Similar result was reported by<sup>25</sup> in Neem extract fed chickens. The significant increase in neutrophil count caused by the fraction 3 probably indicates an enhancement in the ability of the body to attack and destroy invading bacteria, viruses and other injurious agents (phagocytosis). Similar result was reported by<sup>26</sup> in

*Dennettia tripetala* extract treated rats. The significant decrease in the lymphocyte values caused by fraction 3 probably indicates reduction in the acquired immune response of the body. Contrary result was reported by<sup>27</sup> in *Pelargonium reniforme* extract treated rats. Fraction 3 caused significant increase in platelet value which probably indicates an enhancement in the hemostatic function of the body. Similar result was reported by<sup>28</sup> in *Fadogia agrestis* extract treated rats.

Fraction 3 caused significant increase in total protein level which might indicate an enhancement in the buffering capacity of blood and colloid osmotic pressure. Similar result was reported by<sup>27</sup> in *Pelargonium reniforme* extract treated rats. The significant increase in the albumin levels caused by fraction 3 could indicate an increase in the serum levels of metals, ions, fatty acids, amino acids, bilirubin and enzymes; since it has been reported that albumin serves as a carrier for metals, ions, fatty acids, amino acids, bilirubin, enzymes and drugs.<sup>22</sup> Similar result was reported by<sup>29</sup> in *Enicostemma axillare* extract treated rats. Fraction 3 caused significant increase in the globulin levels which probably indicates an enhancement in both the natural and acquired immunity of the body. Similar result was reported by<sup>30</sup> in *Andrographis paniculata* extract treated rats.

Fraction 3 caused significant increase in the activity of ALT, which probably indicates an induction of hepatic

damage, since it has been reported that ALT is present in the liver and other cells and is particularly useful in measuring hepatic necrosis, especially in small animals.<sup>15</sup> Similar result was reported by<sup>31</sup> in *Phyllanthus amarus* extract treated rats. The significant increase in the activity of AST caused by fraction 3 could indicate an induction of tissue necrosis, since it has been reported that elevation in the activity of AST can be associated with cell necrosis of many tissues, pathology involving the skeletal or cardiac muscle and/or hepatic parenchyma, allows leakage of large amounts of this enzyme into the blood.<sup>32</sup> Contrary result was reported by in *Murraya koenigii* extracts treated rats.

## CONCLUSION

In conclusion, this study has shown that chromatographic fraction 3 of *Portulaca oleracea* could have some toxic and beneficial potentialities on the blood chemistry of albino rats. However, its effect on human blood chemistry is unknown; nevertheless, considering these findings in animal model, it is recommended that caution should be exercised in the consumption of *Portulaca oleracea*.

## REFERENCES

- Burkill HM, The useful plants of West Tropical Africa, The White friars Press Limited, Tonbridge, Kent TN9 1QR, Great Britain, vol 4, 1997.
- Johnson, The useful plans of West Africa, The White friars Press Limited, Tonbridge, Kent TN9 1QR, Great Britain, vol 4, 1997.
- Ainslie JR, The list of plants used in native medicine in Nigeria, Imp. Forest. Inst. Oxford Inst., 1937, Paper 7(mimeo).
- Vermeer DE, in litt. dd 28/1/76 re collections ex Benue Plateau and near Benin deposited at Herb UCI, 1976.
- Oyediji K.O, Bolarinwa AF, Adegoke AO, Evaluation of antifertility and teratogenic effects of chromatographic fractions of *Portulaca oleracea* in male and female albino rats, International Journal of Pharmacy and Pharmaceutical Sciences, 5(2), 2013, 691-695.
- Oyediji KO, Bolarinwa AF, Effects of Crude extracts of *Portulaca oleracea* in reproductive functions in male albino rats, IOSR Journal of Pharmacy and Biological Sciences, 4(6), 2013, 71-79.
- Oyediji KO, Bolarinwa AF, Oladosu IA, Effect of isolated improve constituent of *Portulaca oleraceas* on reproductive parameters in male rats. International Journal of Pharmacy and Pharmaceutical Sciences, 5(2), 684-690.
- Miladi – Gorgi H, Vafaei AA, Rashidy – Pour A, Taherian AA, Jarrahi M, Emami-Abargoei M, Investigation of anxiolytic effects of aqueous extract of *Portulaca oleracea* in mice, Iranian Journal of pharmaceutical research, Supplement 2, 2004, 57-57.
- Jain NC, Schalm's Veterinary Haematology 4th ed., 1986, Lea and Fabiger, Philadelphia.
- Dacie JV, Lewis SM, Practical haematology, ELBS with Churchill Livingstone, England, 7, 1991, 37-85.
- Mitruka BM, Rawnsley H, Clinical, biochemical and haematological references values in normal experimental animals, Masson Publishing USA Inc., 1977, 53-54.
- Reinhold JG, Manual determination of serum total protein, albumin and globulin fractions by the Biuret method *Standard Methods of Clinical Chemistry* 1953, Academic Press, New York.
- Dumas BT, Watson W, Biggs HC, Albumin standards and the measurement of serum albumin with bromocresol green, Clinica Chimica Acta, 31, 1971, 87-96.
- Duncan JR, Praise KW, Mahaffey EA, Veterinary Laboratory Medicine (Clinical Pathology) 3rd ed., 1994, Iowa State University Press, U.S.A.
- American Diabetes Association, Nutrition recommendation and principles for people with diabetes mellitus clinical practice recommendations, Diabetes care, 23, 2000, 543-6.
- Olayemi FO, Evaluation of the reproductive and toxic effects of *Cnestis ferruginea* root extract in male rats. Ph.D Thesis, Department of Physiology, 2007, University of Ibadan.
- Adedapo AA, Abatan MO, Olorunsogo OO, Effects of some plants of the spurge family on haematological and biochemical parameters in rats, VETERINARSKI ARHIV, 77 (1), 2007, 29-38.
- Ghai, A textbook of practical physiology, fifth edition. Jaypee Brothers Medical Publishers Ltd, 1999, New Delhi.
- Ibrahim IA, Omer SA, Ibrahim FH, Khalid SA, Adam SE, Experimental *Azadirachta indica* toxicosis in chicks, Veterinary and Human Toxicology, 34, 1992, 221-224.
- Ademola SG, Farinu GO, Oyediji KO, Haematological and serum enzyme activities of broilers fed garlic and ginger supplements, Bowen Journal of Agriculture, Vol. 2(1), 2005.
- Ganong WF, Review of Medical Physiology 22nd edition. McGraw – Hill Companies, 2005, Inc.
- Imoru JO, Eno AE, Unoh FB, Enkanu E, Ofem OE, Ibu JO, Haematopoietic agents in the crude extracts from the leaves of *Viscum album* (mistletoe), Nigerian Journal of Health and Biomedical Sciences, 4(2), 2005, 139-145.
- Guyton AC, Hall J E, Textbook of Medical Physiology 11th edition, 2006, Elsevier Inc.
- Biu AA, Yusufu SD, Rabo JS, Studies of the effects of aqueous leaf extracts of Neem (*Azadirachta indica*) on Haematological parameters in chickens, African Scientist Vol. 10, No 4, 2009.
- Ikpe DE, Nku CO, Effect of ethanolic extract of *Dennettia tripetala* fruit on haematological parameters in albino Wistar rats, Niger J Physiol Sci., 23 (1-2), 2008, 13-17.
- Adewusi EA, Afolayan AJ, Safety evaluation of the extract from the roots of *Pelargonium reniforme* Curtis in male wistar rats, Afr. J. Pharm and Pharmacology, 3(8), 2009, 368-373.
- Yakubu MT, Akanji MA, Oladiji AT, Hematological evaluation in male albino rats following chronic administration of aqueous extract of *Fadogia agrestis* stem, Pharmacognosy Magazine ISSN: 0973-1296, 2007.



28. Gite VN, Pokharkar RD, Chopade VV, Takate SB, Hepato-Protective activity of *Enicostemma Axillare* in paracetamol induced hepato-toxicity in albino rats. International Journal of Pharmacy and Life Sciences, www.ijplsjournal.com 2010.
29. Puri A, Saxena R, Saxena RP, Saxena KC, Immunostimulant agents from *Andrographis paniculata*, J Nat Prod, 56, 1993, 995-999.
30. Taiwo IA, Oboh BO, Francis-Garuba PN, Haematological properties of aqueous extracts of *Phyllanthus amarus* and *Xylopiia aethiopica* A. Rich in albino rats, Ethno-Med, 3(2), 2009, 99-103.
31. Bush BM, Interpretations of Laboratory Results for Small Animal Clinicians. Blackwell Scientific Publications, 1991, London.
32. Adebajo AC, Ayoola OF, Iwalewa EO *et al*, Anti-trichomonal, biochemical and toxicological activities of methanolic extract and some carbazole alkaloid isolated from the leaves of *Murraya koenigii* in Nigeria, Phytomedicine, 13, 2006, 246-254.

<b>Source of Support: Nil, Conflict of Interest: None.</b>
--