

# Magnesium Deficiency and its Correlation with Insulin Resistance in Obese Females in Majmaah

\*Johra Khan

\*Assistant Professor, MDL Dept., College of Applied Medical Sciences (female), Majmaah University, Al- Majmaah, 11952, Saudi Arabia. \*Corresponding author's E-mail: j.khan@mu.edu.sa

Received: 10-03-2018; Revised: 28-04-2018; Accepted: 12-04-2018.

### ABSTRACT

The aim of this study is deficiency of Magnesium is related to development of insulin resistance that later increases the risk of type 2 diabetes. This study was designed to find whether serum magnesium is exhibited in obese female and its relation with insulin resistance. We selected 50 non-diabetic females (BMI> 30) calculated their BMI and measured serum magnesium and glucose using kits. Serum magnesium was found significantly lower in obese females (0.640 + 0.015 mmol/I, means + SE) compared with the lean control females (0.821 + 0.012 mmol/I). It was found that serum magnesium was inversely correlated with insulin (rs = -0.36 [95% Cl -0.59 to -0.08]; P = 0.011). The association between magnesium deficiency and insulin resistance is present in obese females. Serum magnesium deficiency in obese females may be secondary to decreased dietary magnesium intake. Magnesium supplementation or increased intake of magnesium- rich foods may be an important tool in the prevention of type 2 diabetes in obese females.

Keywords: Magnesium, Deficiency, Obesity, Insulin resistance.

#### **INTRODUCTION**

besity is often defined as multi-faceted disorder associated with an increase in number of diseases like hypertension, diabetes mellitus, arthritis, gout, coronary heart diseases and gall bladder <sup>1, 2</sup>. The rapid social and cultural change that has taken place in Arabian and Gulf countries after the discovery of oil during 1970s and 1980s increased obesity at an alarming rate in these countries <sup>3, 4, 5</sup>. Many studies show that obesity is a major risk factor for chronic diseases and plays an important role in the "insulin resistance" or "metabolic" syndrome, which includes much disease like hyperinsulinemia, hypertension, hyperlipidemia, type 2 diabetes mellitus, and an increased risk of atherosclerotic cardiovascular disease. The metabolic syndrome is a common pathophysiologic condition with implications for the development of many chronic diseases. Obesity childhood often beginning in precedes the hyperinsulinemia state <sup>15</sup>. The developing countries are observing increasing incidence of childhood obesity as a result of this new cases of metabolic syndrome among children are increasing, which in turn is likely to create an enormous socioeconomic and public health burden for nations with large number of young population like KSA in the near future<sup>16</sup>. However, little is known about the prevalence of childhood obesity and the pediatric metabolic syndrome because of the limited number of studies, the various definitions used, and the different age groups studied, which makes comparisons difficult. Many studies conducted in kingdom of Saudi Arabia (KSA) revealed that a major change in dietary habits has taken place. Fat consumption and fast food consumption increased as well as processed food has become a major part of everyday meal <sup>6</sup>.

Magnesium is fourth abundant material in body and an important cofactor for enzymes involved in Carbohydrate metabolism. Many studies show a strong relation between magnesium and insulin action and a low serum and intercellular magnesium are associated with insulin resistance <sup>7</sup> impaired glucose tolerances and decreased insulin secretion <sup>8</sup>. The present study was designed to study whether serum magnesium is exhibited in obese and its relation with insulin resistance.

## MATERIALS AND METHODS

We selected 30 non-diabetic females (BMI>25) age between 20 to 25 years. 20 lean females (BMI<25) matched for sex and stage of pubertal development. Female students were recruited with written informed consent obtained from both parents and students. On arrival, a questionnaire was provided to fill all necessary information and weight was measured using a calibrated digital scale. Height was measured in triplicate to the nearest millimeter using a calibrated stadiometer. BMI scores were calculated using equation provided by Centre for diseases control and prevention (CDC).

#### Laboratory Tests

Blood samples were obtained for glucose and serum magnesium. Serum glucose were measured using glucose liquicolour kit (Human) which is an enzymatic colorimetric method. The optimum density (OD) was measured at 500nm. Serum magnesium was measured using magnesium measuring kit (Human) which is also enzymatic colorimetric method. The optimum density (OD) was measured at 470nm.



International Journal of Pharmaceutical Sciences Review and Research Available online at www.globalresearchonline.net 117

© Copyright protected. Unauthorised republication, reproduction, distribution, dissemination and copying of this document in whole or in part is strictly prohibited.

## Statistical analysis

Difference between obese and lean female students were examined for serum glucose and serum magnesium. *P* values for these comparisons were calculated via ANOVA. Weight status (lean, obese) served as independent factors of the ANOVA model. Statistical significance was set at  $\alpha$  < 0.05.

# **RESULTS AND DISCUSSION**

# Serum Magnesium

Serum magnesium concentration was significantly lower in obese females compared with lean females 0.472  $\pm$  0.015 vs lean 0.903  $\pm$  0.012 mmol/l; *P* = 0.02.

# Serum Glucose

Serum glucose concentration was significantly high in obese females compared with lean females  $0.726 \pm 0.011$  vs lean  $0.387 \pm 0.015$  mmol/l; P = 0.03.

**Table 1:** Characteristics and metabolic parameters of the study subjects

Parameter	Obese females	Lean females	*P values
n (number of subjects)	30	20	
BMI (kg/m²)	30.5 <u>+</u> 2.1; 28.4 (24.5 – 35)	22.5 <u>+</u> 0.6; 21.4 (20 – 24.5)	<0.001
Fasting glucose (mmol/l)	5.80 <u>+</u> 0.11; 5.69 (5.6 – 6.8)	5.16 <u>+</u> 0.015; 5.15 (4.8 – 5.5)	0.782
Serum magnesium(mmol/I)	0.472 <u>+</u> 0.015; 0.467 (0.402 – 0.653)	0.93 <u>+</u> 0.012; 0.891 (0.732 – 1.01)	0.005

\*By ANOVA for continuous variables and by exact test for categorical variables.

### Serum Magnesium and Glucose

Serum magnesium was found inversely proportional to fasting serum glucose level in obese female students.

# CONCLUSION

Our studies found that low serum magnesium concentrations are associated with hyperinsulinemia, decreased insulin-mediated glucose disposal, and the metabolic syndrome<sup>9</sup>. In the Atherosclerosis Risk in Communities study, low serum magnesium had a two fold increase in incidence of type 2 diabetes compared with those having a normal magnesium concentration<sup>10</sup>. Magnesium supplementation in subjects with IR and type 2 diabetes resulted in improvement of insulin sensitivity<sup>11</sup> and cell response to glucose<sup>12</sup>. In an animal model of the metabolic syndrome, increased magnesium intake reduced the rate of development of type 2 diabetes<sup>13, 14</sup>. These results suggest a potential role of magnesium in the prevention of type 2 diabetes. It will be important to test in future interventional studies whether increasing magnesium intake will improve insulin sensitivity and reduce the risk of type 2 diabetes in obese females.

# REFERENCES

- 1. World Health Organization: Obesity: preventing and managing the global epidemic. Report of a WHO consultation. World Health Organ Tech Rep Ser. 894, 2000, 1 -253.
- James WPT, Jackson Leach R, Mhurchu CN, Kalamara E, Shayeghi M, Rigby NJ, Nishida C, Rodgers A: Overweight and obesity (high body mass index). Comparative quantification of health risks: global and regional burden of disease

attributable to selected major risk factors. Edited by: Ezzati M, Lopez AD, Rodgers A, Murray CJL. 2004, 497- 596.

- Yang P, Zhou Y, Chen B, Wan HW, Jia GQ, Bai HL, Wu XT: Overweight, obesity and gastric cancer risk: Results from a meta-analysis of cohort studies. Eur J Cancer. 45 (16), 2009, 2867-2873.
- DeFronzo RA, Tobin JD, Andres R: Glucose clamp technique: a method for quantifying insulin secretion and resistance. Am J Physiol 237, E214 – E223, 1979.
- 5. Laakso M: How good a marker is insulin level for insulin resistance, Am J Epidemiol 137, 959 965.
- Matthews DR. Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC: Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentration in man. Diabetologia 28, 1985, 412 – 419.
- Legro RS, Finegood D, Dunaif A: A fasting glucose to insulin ratio is a useful measure of insulin sensitivity in women with polycystic ovary syndrome. J Clin Endocrinol Metab, 83, 1998, 2694 – 2698.
- Emoto M, Nishizawa Y, Maekawa K, Hiura Y, Kanda H, Kawagishi T, Shoji T, Okuno Y, Morii H: Homeostasis model assessment as a clinical index of insulin resistance in type 2 diabetic patients treated with sulfonylureas. Diabetes Care 22, 1999, 818 – 822.
- Bonora E, Targher G, Alberiche M, Bonadonna RC, Saggiani F, Zenere MB, Monauni T, Muggeo M: Homeostasis model assessment closely mirrors the glucose clamp technique in the assessment of insulin sensitivity: studies in subjects with various degrees of glucose tolerance and insulin sensitivity. Diabetes Care 23, 2000, 57 – 63.



Available online at www.globalresearchonline.net

© Copyright protected. Unauthorised republication, reproduction, distribution, dissemination and copying of this document in whole or in part is strictly prohibited.

- 10. Hermans MP, Levy JC, Morris RJ, Turner RC: Comparisons of insulin sensitivity tests across a range of glucose tolerance from normal to diabetes. Diabetologia 42, 1999, 678 687.
- 11. Katz A, Nambi SS, Mather K, Baron AD, Follmann DA, Sullivan G, Quon MJ: Quantitative insulin sensitivity check index: a simple, accurate method for assessing insulin sensitivity in humans. J Clin Edinocrinol Metab 85, 2000, 2402 2410.
- Mather KJ, Hunt AE, Steinberg HO, Paradisi G, Hook G, Katz A, Quon MJ, Baron AD: Repeatability characteristics of simple indices of insulin resistance: implications for research applications. J Clin Endocrinol Metab 86, 2001, 5457 – 5464.
- Perseghin G, Caumo A, Caloni M, Testolin G, Luzi L: Incorporation of the fasting plasma FFA concentration into QUICKI improves its association with insulin sensitivity in

non-obese individuals. J Clin Endocrinol Metab 86, 2001, 4776 – 4781.

- 14. Paolisso G, Passariello N, Pizza G, Marrazzo G, Giunta R, Sgambato S, Varricchio M, D'Onofrio F: Dietary magnesium supplements improve B-cell response to glucose and arginine in elderly non-insulin dependent diabetic subjects. Acta Endocrinol Copenhagen 121, 1989, 16-20.
- 15. Van Dam RM, Rimm EB, Willett WC, Stampfer MJ, Hu FB. Dietary patterns and risk for type 2 diabetes mellitus in U.S. men. Ann Intern Med 136, 2002, 201–9.
- 16. Schulze MB, Hu FB. Primary prevention of diabetes: what can be done and how much can be prevented? Annu Rev Public Health 26, 2005, 445–67.

Source of Support: Nil, Conflict of Interest: None.

