

Research Article



Combined *in vivo* Antioxidant Activity of Crude Seed Extracts of Selected Medicinal Plants on High Fat Diet Induced Rats

Zafar Javed Khan^{1*}, Naeem Ahmad Khan¹, Imrana Naseem², Shahab A A Nami³

¹Department, of Ilmul Advia, Faculty of Unani Medicine, Aligarh Muslim University, Aligarh, UP, India.

²Department of Biochemistry, Faculty of Life Science, Aligarh Muslim University, Aligarh, UP, India.

³Department of Kulliyat, Faculty of Unani Medicine Aligarh Muslim University, Aligarh, UP, India.

*Corresponding author's E-mail: zafarjaved9454@gmail.com

Received: 02-08-2019; Revised: 24-09-2019; Accepted: 30-09-2019.

ABSTRACT

The objective of the study is to evaluate the *in-vivo* antioxidant potency of 50% ethanolic extract of seeds of *Linum usitatissimum*, *sesamum indicum*, and *Moringa oleifera* fruits extract against high fat diet induced rats, when used in combinations. Animal were treated with compound extract for 30 days, and high fat diet was given to all groups except plain control through, out the study, and alpha tocopherol acetate (Vit, E) was used as standard. Pre-treatment with 22.4 mg/100 gm of body weight of 50% ethanolic extract of compound extract improved the Superoxide dismutase, catalase, glutathione, and lipid peroxidation levels significantly as compared to control group. The present studies revealed that combined extract has significant *in-vivo* antioxidant activity and can be use to protect tissue from oxidative stress. The result showed that the activities of SOD, catalase, lipid peroxidase, and glutathione, in groups treated with high fat diet declined significantly than that of normal group. 50% ethanolic extract of compound in the dose of 22.4 mg/100 gm of body weight, has improved the SOD, catalase, glutathione, and lipid peroxidation levels significantly, as compared to the alpha tocopherol (Vit,E). Based on this study we conclude that 50% ethanolic extract of compound possesses *in vivo* antioxidant activity and can be employed in protecting tissue from oxidative stress.

Keywords: Compound extract, alpha tocopherol, superoxide dismutase, antioxidant activity.

INTRODUCTION

The mixture of bioactive constituents and their byproducts contained in plant extracts and various natural products produce synergistic effects. But the mechanism responsible for synergistic antioxidant activity, are not explained yet due to the complex nature of the plant extracts. The synergistic antioxidant activity of the optimized mixture can more effectively or at lower doses prevent or treat the various diseases caused by oxidative damage. The present study was conducted on a non-pharmacopoeial compound extract (NPCE). This is a combination of three drugs viz. Tukhm-e-Katan comprises of seeds of a plant *Linum usitatissimum* Linn. Versatile and blue flowering rabi crop belonging to linaceae family, commonly known as Flaxseed or Linseed¹. The plant has shown diverse biological and pharmacological activities. It has been used in Unani medicine and traditional systems of medicine from time immemorial. Its seed and oil are used in various diseases such as asthma, cough, bronchitis, pleurisy, pneumonia, joint pain, renal colic, renal calculi, rheumatic swelling, Katan is a famous Unani drug used in a number of pathological conditions. Although entire plant has medicinal value but its seed and oil are more important and have broad medicinal values. Katan is an annual herb of about 0.7 m high with blue flowers and a globular capsule. The seed are ovate, flattened and obliquely pointed at one end, about 4-6 mm long and 2-2.5 mm broad². According to Unani it has Mohallil-e-waram^{3,4}, Dafe-e-Sua'al, Muqawwi-e-Bah Muqawwi-e-Aaza and Mulayyin, properties^{3,5}. Sehjana comprises of Fruits/Podsof *Moringa oleifera* Lam,

(Moringaceae) commonly known as Sehjana in Unani Medicine. The seed of pods (fruits) of the *Moringa oleifera* tree are one of the most nutritive and useful parts of this versatile plant. Various parts of this plant act as cardiac and circulatory stimulants, possess antitumour, antipyretic, antiepileptic, anti-inflammatory, antiulcer, antispasmodic, diuretic⁶, antihypertensive, cholesterol lowering⁷, and antioxidant, antidiabetic, hepatoprotective, antibacterial and antifungal activities^{8,9}, aphrodisiac, anthelmintic, analgesic activities^{10,11}. According to Unani literature it possesses many actions like Moohallil-e-Waram, Muqawwi-e-Bah, Mushtahi, Qatile Kiram-e-Amaa^{3,12}. So medicinally used in Waja-ul-Mafasil, Waja-ul-Qutu, Zof-e-Ishteha^{12,13}. And Kunjad sufaid consists of dried seeds of *sesamum indicum* Linn, commonly known as Til, sesame or benne seed, is cultivated through, out India, mainly for its seeds and oil, in India, due to the presence of bioactive components present in the seed including polyunsaturated fatty acids, phytosterols, tocopherols, vital minerals and unique class of phenylpropanoid compounds namely lignans such as sesamin, sesamol and sesamol¹⁴. These phytochemicals provide defense mechanism against reactive oxygen species and increases keeping quality of oil by preventing oxidative rancidity^{15,16}. Sesame lignans have various pharmacological properties including, antioxidant activity^{17,18}, antimicrobial activity¹⁹ anti-proliferative activity²⁰ lowering cholesterol levels²¹. Sesame seed have been used as a medicine since antiquity. They are considers to act as aphrodisiac, demulcent, lactagogue, emmenagogue, diuretic, and laxative^{8,10,7}. The main aim



of the study is to provide insight about the combined Antioxidant activity of Crude Seed extracts of selected medicinal plants on High Fat Diet Induced Rats.

MATERIAL AND METHODS

Collection of plant

The seeds of *Linum usitatissimum* and *Sesamum indicum* were procured from local market of Aligarh. Where the pods/ fruits of *Sehjana (Moringa oleifera Lam.)* was collect directly from the herbal garden of department of Ilmul Advia AMU, Aligarh, and are properly identified according to the botanical, Unani and Ayurvedic literature and then confirmed in pharmacognosy section of department of Ilmul Advia. A herbarium sample of the test drugs were prepared and submitted to mawalid-e-salasa museum of the department after identification for further reference.

Preparation of extracts

The test drugs specially *Moringa* was cleaned from the earthy material, washed with double distilled water and shade dried to powdered in electrical grinder with slow and light movement to avoid sticking of the drug material with the grinder and there after the drug was passed through the sieve no. 80 to confirm its fineness and uniformity of particle size. And the powder was packed in to soxhlet apparatus and extracted with 50% ethanol (64.5 – 65.5°C) The extract was filtered and concentrated by evaporation on water bath, the yield percentage was calculated with reference to crude drug.

Preliminary Physicochemical and phytochemical screening

The physicochemical study of the test drugs included the study for organoleptic characters, ash value, moisture content, pH value, loss of weight on drying, successive extractive value, alcohol and water soluble, matter, bulk density. And the phytochemical screening was carried out with different extract of compound extract for the detection of various phytochemicals. Tests for common phytochemicals were carried out by standard methods²².

Drugs

The test drugs were obtained as describe above, sample of all drugs was found to be the standard in light of our physicochemical studies, therefore, they used for pharmacological studies. α -tocopherol acetate, used as the standard drug, was obtained from Loba chemical. The hydro-alcoholic (50% alcohol and 50% water) extract of drug was use for the study. The yield percentage was calculated with reference to dried drug. The extract of test drugs was dissolved in distilled water the dose was obtained by multiplying the Unani clinical dose with appropriate conversion factor of 7 for rats²³, and found to be 15 mg/kg BW for α -tocopherol acetate, and 22.4 mg/100gm BW for compound extract. A feeding canula was use to administer the suspension, which was homogenized by shaking well for two minutes.

Diet

Animals during the acclimation period were fed the commercially available rat chow (Ashirwad diet). Afterwards a special high lipid diet/atherogenic diet²⁴, containing butter (5g), bread slice (1), wheat flour 3 tea spoon), milk powder (1.5 tea spoon), cholesterol powder (60 mg/kg) and coconut oil (1 ml) was also given during experimentation period along with the normal diet to all the animals except those in plain control. The high lipid diet was given every day in the morning to the animals and when they consumed it they were allowed free access to the normal diet.

Animals

Male albino rats (Wister strain) of same age, weighing 150-200 gm, were purchase from the central animal house of Indian veterinary research institute Bareilly, Up, India. The animals were housed in sufficiently, large cages and treated under humane and hygienic condition with maintained at uniform temperature $25 \pm 2^\circ\text{C}$ and 12 h day: night cycle according to departmental ethical committee for animal experimentation. and were fed on standard diet (Ashirwad industries, Chandigarh, India) and tape water, *ad libitum*. The animal is deprived of food for 12 hours before the administration of treatment, water was provided throughout the study. The animals in all the, group were administered with the treatment by oral route once a day for 30 days. Before starting the experiment, permission from the Institutional Animal Ethics Committee was obtained, animal experimentations were permitted by Ministry of environment and forests, government of India under registration no, 714/02/aCPCSEA. It was issue by committee for the purpose, of control and supervision of experiments on animals (CPCSEA) dated 15th September 2016 and approved by the Institutional Animal Ethics Committee (IAEC) of department of biochemistry, Faculty of Life Science, Aligarh Muslim University, Aligarh, India.

Experimental design

Rats were divided randomly into five group of six animals each and treated for 30 days as follows. Group-I animals served as plain control, treated with vehicle (vehicle control). Group-II served as control group treated with high fat diet and normal diet throughout the study. Group –III served as a standard group, and was administered alpha tocopherol acetate in a dose of 15 mg/kg bw. Group-IV was treated with normal diet and with daily dose of 22.4 gm/100 gm,bw. of ethanolic extract of compound extract. Group-V was treated with high fat diet and with daily dose of 22.4 gm/100 gm,bw. of ethanolic extract of compound extract for 30 days. High fat diet was given in all groups except in plain control group and group treated with normal diet. 6 h after the last treatment, on day 31 the rats were anaesthetized by chloroform and sacrificed, all efforts were made to minimize suffering, blood, liver, and brain was rapidly excised, rinsed in ice-cold saline, and a 10% w/v homogenate was prepared



using 0.15M KCl, centrifuged at 800 rpm for 10 min at 4°C. The supernatant obtained was used for the estimation of catalase, lipid peroxidase, and other enzymes. Further, the homogenate was centrifuged at 1000 rpm for 20 min at 4°C and the supernatant was used for biochemical estimation.

Collection and preparation of biological samples

After sacrificing the animals, the blood was collected and centrifuged at 2500 x g for 10 minutes at 4°C and the separated serum was collected carefully. The liver and brain were, also removed immediately washed with chilled normal saline and preserved in ice. A 10% W/V homogenate was prepared in chilled 0.15M KCL for Lipid Peroxidation, and Superoxide Dismutase; in 0.1M Chilled Tris HCl buffer (pH 8.2) for Glutathione Reductase, and 50mM phosphate buffer for Catalase. The parameter estimated in serum, liver and brain were Lipid peroxidation, Superoxide Dismutase, Catalase, and Glutathione Reductase.

Statistical Analysis

The concentration of each parameter in various animal groups (Gp I- V) were statistically compared for determining significance of difference by one-way ANOVA test followed by pair-wise comparison of various groups by LSD. P value of 0.05 or less was considered significant. The analysis was carried out by using the software of the website, www.myassay.com. Values are presented as mean ± standard deviation for groups of six animals.

Biochemical estimation

Estimation of superoxide dismutase (SOD)

The activity of superoxide dismutase was estimated by Elisa reader with the help of commercially available Detectx Superoxide Dismutase (SOD) colorimetric activity kit. (K028-H1) Arbor Assays USA.

Principle

The substrate is added followed by Xanthine oxidase reagent and incubated at room temperature for 20 minutes. The Xanthine oxidase generates superoxide in the presence of oxygen, which convert a colorless substrate in the detection reagent into a yellow colored product. The colored product is read at 450 nm. Increasing levels of SOD in the samples causes a decrease in superoxide concentration and reduction in yellow product. The activity of the SOD in the sample is calculated after making a suitable correction for any dilution, and expressed in terms of unit of SOD activity per mL

Estimation of catalase

The catalase activity was determined by elisa reader with the help of commercially available Detectx Catalase colorimetric activity kit. (K033-H1) Arbor Assays USA.

Principle

Samples are diluted in the provided assay, buffer and added to the wells of a half area clear plate. Hydrogen peroxide is added to each well and the plate incubated at room temperature for 30 minute. The supplied substrate is added, followed by diluted horseradish peroxidase and incubated at room temperature for 15 minutes. The HRP reacts with the substrate in the presence of hydrogen peroxide to convert the colorless substrate into a pink-colored product. The colored product is read at 560 nm. Increasing levels of catalase in the samples causes a decrease in H₂O₂ concentration and reduction in pink product, and expressed in terms of units of Catalase activity per mL.

Estimation of lipid peroxidation (TBARS)

Lipid peroxidation was estimated by Elisa reader with the help of commercially available Quantichrom™ TBARS Assay kit (DTBA-100), Bioassay Systems USA.

Principle

Bioassay system' TBARS assay is based on the reaction of TBARS with thiobarbitturic acid (TBA) to form a pink colored product. The color intensity at 535 nm or fluorescence intensity at (λ_{ex}/em =560 nm / 585 nm) is directly proportional to TBARS concentration in the sample, and expressed in terms of μM MDA (μM=μmole/liter=nmole/ml).

Estimation of glutathione reductase (GR)

The Glutathione reductase activity was estimated by elisa reader with the help of commercially available Quantichrom™ Glutathione Reductase Kit (ECGR-100), Bioassay Systems USA.

Principle

Bioassay system' non-radioactive, colorimetric GR assay is designed to accurately measure GR activity in biological samples with a method that utilizes Ellman's method in which DTNB reacts with the GSH generated from the reduction of GSSG by the GR in a sample to form a yellow product (TNB²⁻). The rate of changes in the optical density, measured at 412 nm, is directly proportional to GR activity in the sample, and expressed in terms of unit per mL, the 1 Unit (U) of GR will catalyze the conversion of 1 μmole of GSSG to 2 μmole GSH per min at pH 7.6.

Estimation of lipid profile

Estimation of Cholesterol

Cholesterol was estimated by photocolormeter with the help of commercially available cholesterol (SR) kit based on CHOD / PAP method, (Erba Mannheim Germany).

Principle

The estimation of cholesterol involves the following enzyme catalyzed reactions.









Estimation of Triglyceride

Triglyceride was estimated by photocolorimeter with the help of commercially available triglyceride (SR) kit based on GPO / PAP method, (Erba Mannheim Germany).

Principle




$$\text{Triglyceride (mg/dL)} = \frac{\text{Abs.of Test}}{\text{Abs.of Standard}} \times \text{concentration of standard (mg/dL)}$$

Estimation of HDL Cholesterol

HDL cholesterol was estimated by photocolorimeter with the help of commercially available HDL cholesterol PPT. set based on phosphotungstic acid method, (Erba Mannheim Germany).

Principle

chylomicrons, LDL and VLDL (low and very low density lipoproteins) are precipitated from serum by phosphotungstate in the presence of divalent cations such as magnesium. The HDL cholesterol remain unaffected in the supernat and is estimated using ERBA cholesterol reagent.



$$\text{HDL Cholesterol (mg/dl)} = \frac{\text{Abs.of Test}}{\text{Abs.of Std.}} \times \text{concentration of standard (mg/dL)} \times \text{dilution factor}$$

$$= \frac{\text{Abs.of Test}}{\text{Abs.of Standard}} \times 25 \times 3$$

$$= \frac{\text{Abs.of Test}}{\text{Abs.of Standard}} \times 75$$

Estimation of LDL and VLDL

The values of LDL were calculated by following formulae.

LDL = Total cholesterol – HDL – VLDL (Friedewald formulae) VLDL = Triglyceride / 5 (Friedewald WI, 1972). The atherogenic index of plasma was calculated by the formula AIP = [TGL/HDL]. While HDL/LDL ratio by dividing the value of HDL with that of LDL.

RESULTS

The present study determines a comprehensive range of physicochemical characters of the drug according to the parameters used in pharmacopeia, which may serve as

the standard for ensuring optimum efficacy and safety of various samples of the drug.

Phytochemicals investigation

It was founding that different extract of compound contained alkaloids, phenol, carbohydrate, protein, amino acids, steroids, glycosides and tannins.

In-vivo antioxidant activity

The present study was undertaken to assess the in-vivo antioxidant potential of 50% ethanolic extract of compound drug against high fat diet induced rats, in serum, liver, and brain homogenate of control and experimental groups of rats. The results showed that the activities of superoxide dismutase (SOD), catalase (CAT), and glutathione reductase (GR) in the liver, brain and serum of control and experimental groups of rats was significantly lower in the high fat diet control group as compared to the plain control group. In the standard group, and compound treated group the activity of these enzymes was significantly, increased in comparison to that in the control group. (Table 1 to 3).

Table 4 shows the activities of lipid peroxidation (TBARS) in the liver, brain and serum of control and experimental groups of rats. The activity of lipid peroxidation (TBARS) in liver, brain and serum was significantly elevated in the high fat diet control group as compared to the plain control group. In the standard group, and compound treated group the activity of (TBARS) was significantly lower in comparison to that in the control group. (Table 4).

Table 5 shows the activities of lipid Profile in serum of control and experimental groups of rats. The activity of lipid profile in serum was significantly elevated in the high fat diet control group as compared to the plain control group. In the standard group, and compound treated group the activity of lipid profile was significantly lower in comparison to that in the control group. (Table 5).

The test drugs exhibit high efficacious antioxidant activity. They are shown to be more effective than the standard antioxidant agents, the test drugs shown to be comprehensive antioxidant agents as they have been found to be effective in three biological samples, namely liver and brain homogenate and serum.




Figure 1: Activity of superoxide dismutase (SOD) in serum, liver and brain (U/mL).



Table 1: Effect of 50% ethanolic extract of combined extract on the activity of Superoxide dismutase (SOD) in high fat diet induced rats, in serum, liver and brain.


Superoxide dismutase (SOD) (U/mL)*			
Groups	Serum	Liver	Brain
Plain control	1.05±0.02	2.81±0.06	1.71±0.31
Control	0.69±0.05 b***	1.79±0.02 b***	1.08±0.00 b**
Standard	1.10±0.07 a***b*	2.95±0.03 a***b*	2.17±0.02 a***b*
Combination ē normal diet	1.01±0.01 a**	2.54±0.02 a***	1.95±0.03 a***
Combination ē high fat diet	0.98±0.00 a**	2.43±0.04 a***	1.68±0.04 a**

(n=6); Values are in Mean ±SEM. Where* P<0.05 and **P<0.01 *** p<0.001; a = Against control, b = Against plain control, c = Against standard; * The results are expressed in term of unit of SOD activity per mL.

Table 2: Effect of 50% ethanolic extract of combined extract on the activity of catalase in high fat diet induced rats, in serum, liver and brain.


Catalase (U/mL)*			
Groups	Serum	Liver	Brain
Plain control	1.18±0.019	3.41±0.037	1.57±0.029
Control	0.97±0.020 a***	2.22±0.059 a***	1.16±0.025 a***
Standard	1.32±0.031 a*** b**	3.92±0.039 a***b***	2.00±0.013 a***b***
Combination ē normal diet	1.29±0.031 a***	3.11±0.033 a*** b***	1.73±0.044 a*** b**
Combination ē high fat diet	1.09±0.022 a*	3.02±0.035 a*** b***	1.43±0.020 a*** b**

* The results are expressed in term of units of Catalase activity per mL.

**Figure 2:** catalase activity in serum, liver and brain (U/mL)**Table 3:** Effect of 50% ethanolic extract of combined extract on the activity of Glutathione Reductase (GR) in high fat diet induced rats, in serum, liver and brain.

Glutathione reductase (U/L)*			
Groups	Serum	Liver	Brain
Plain control	2.73 ±0.061	4.37 ±0.153	3.64±0.066
Control	1.66±0.061 a***	3.22 ±0.147 a***	2.60±0.148 a***
Standard	3.21±0.053 a*** b***	5.38 ±0.188 a***b***	4.44 ±0.232 a*** b***
Combination ē normal diet	3.70±0.072 a*** b*** c***	4.54 ±0.072 a*** c***	3.96±0.037 a*** c**
Combination ē high fat diet	3.31±0.070 a*** b***	4.43±0.065 a*** c***	3.78 ±0.074 a*** c**

* The 1 Unit (U) of GR will catalyze the conversion of 1 μmole of GSSG to 2 μmole GSH per min at pH 7.6.

**Figure 3:** Glutathione reductase (GR) activity in serum, liver and brain (U/mL).**Table 4:** Effect of 50% ethanolic extract of combined extract on the activity of Lipid Peroxidation in high fat diet induced rats, in serum, liver and brain.

Lipid peroxidation (TBARS) μM MDA (μM=μmole/liter=nmole/ml)			
Groups	Serum	Liver	Brain
Plain control	3.05±0.03	5.28±0.05	5.87±0.06
Control	4.63±0.06 b***	6.68±0.11b***	7.96±0.03 b***
Standard	2.65±0.12 a*** b**	4.38±0.08 (a,b)***	5.47±0.09 (a,b)***
Combination ē normal diet	3.17±0.01 a***	5.45±0.09 a***	6.05±0.03 a*** x**
Combination ē high fat diet	3.31±0.01a***	5.72±0.11a***	6.24±0.04 a***




Figure 4: Lipid peroxidation (TBARS) in serum, liver and brain (µM MDA).

Table 5: Effect of combined extract on Lipid Profile in high fat diet induced Rats (Mean±SE).

Group	Lipid profile (mg/dl)						Atherogenic Index of Plasma
	Total Cholesterol mg/dl	Triglyceride (mg/dl)	HDL (mg/dl)	LDL (mg/dl)	VLDL (mg/dl)	HDL:LDL	
Group I P. control	90.87±0.98	159.18 ±2.69	28.2 ±0.77	33.74 ±1.20	30.75 ±0.56	0.83 ±0.02	0.751
Group II Control	104.74±0.66 a***	185.67 ±2.45 a***	24.11 ±1.13 a***	44.59 ±1.19 a*	36.07 ±0.79 a***	0.53 ±0.01	0.886
Group III Standard	96.12±1.67 a*	173.18 ±2.47 a*** b***	28.73 ±0.55 a*** b***	32.62 ±1.87 a**	34.63 ±0.40 b***	0.89 ±0.05	0.780
Group IV	79.03±2.46 a***	167.35 ±1.04 a*** b**	30.55 ±0.59 a*** b**c***	15.22 ±2.53 (a,b,c)***	33.46 ±0.20 a**b***	2.12 ±0.26 a*** b* c***	0.738
Group V	81.25±1.53 a***	168.6 ±2.19 a*** b**	30.05 ±0.52 a*** b**c***	17.50 ±1.64 (a,b,c)***	33.67 ±0.43 a*b***	1.81 ±0.21a*** b**c*	0.749




Figure 5: Effects of test drugs on Lipid profile.

DISCUSSION

The present study, was evaluate the potential effects of 50% ethanolic extract of combination on antioxidant status in high fat diet induced rats. The above findings show that compound extract produce a striking increase in the activity of SOD in all the 3 samples studied viz liver,

brain and the serum, which is greater than the SOD activity in the standard group and even that in the plain control group. Superoxides dismutase is one of the most early and fundamental means of combating the excess ROS, by converting the superoxide ion into the relatively, less reactive oxygen and hydrogen peroxides and thus form a crucial part of the cellular antioxidant defense



mechanism²⁵, $2O_2^{\cdot-} + 2H^+ + SOD \rightarrow H_2O_2 + O_2$ the superoxide ion is itself reactive and may cause oxidative damage, but its harmfulness lies mainly in its being the progenitor of the more reactive and dangerous hydroxyl radical²⁶. Therefore, the striking increase in SOD activity indicates that the test drugs oppose the oxidative damage at an early and crucial point by preventing the generation of one of the most reactive and dangerous oxidative group. Catalase is a ubiquitous antioxidant enzyme that is present in most aerobic cells. Catalase is involved in the detoxification of hydrogen peroxide (H_2O_2), a reactive oxygen species (ROS), which is a toxic product of both normal aerobic metabolism and pathogenic ROS production. The catalase completes the task initiated by SOD by converting less reactive H_2O_2 , produced by SOD, into two molecules of water and harmless molecular oxygen²⁷. The test drugs produce a striking increase in catalase activity along with that in SOD activity, show that they very effectively prevent the generation of ROS and hence, the antioxidant activity is likely to be strong, comprehensive and complete, due to effective prevention and blockade of ROS generation will protect all biomolecules. Thus, the present study shows that MDA concentration is significantly, increased in the high fat diet control group. Malondialdehyde (MDA) is one of the many products of lipid peroxidation caused by reactive oxygen species (ROS), therefore, the increase in MDA concentration indicate an increase in Lipid peroxidation. Malondialdehyde (MDA) is a naturally occurring product of lipid peroxidation, lipid peroxidation is a well-established mechanism of cellular injury in both plants and animals and is used as an indicator of oxidative stress in cells and tissues^{28,29}. During lipid oxidation, malanoaldehyde (MDA) can react with the free amino group of proteins, phospholipids, and nucleic acids damaging their structure and functions. Increased levels of lipid oxidation products are associated with diabetes and atherosclerosis^{30,31}. Oxidative stress, i.e. induced generation of ROS that cannot be fully antagonized, by physiological antioxidants, results in oxidative damage to all biomolecules, however, lipid damage is the most important and takes the form of lipid peroxidation. The significant decrease in MDA concentration shown in the standard group, administered with standard antioxidant agent Vit E (α -tocopherol acetate), indicate the integrity and validity of our experimental procedure by showing the expected protective effect of Vit E against Lipid peroxidation. So the increase in lipid peroxidation and decreases in SOD, Catalase and GR concentration shows that the high fat diet induced in rats successfully, causes oxidative stress and damage. Therefore, lipid rich diets also capable of generating ROS because of antioxidant enzymes they can alter oxygen metabolism. Upon the increase of adipose tissue, the activity of antioxidant enzymes, such as SOD, CAT and glutathione peroxidases (GPx), was found to be significantly diminished³², finally, high ROS production and the decrease in antioxidant capacity, leads to various abnormalities especially, endothelial dysfunction, in a study³³, showed that a diet

high in fat and carbohydrates induces a significant increase in oxidative stress and inflammation in person with obesity. Therefore, we observed a significant reduction in antioxidant enzymes such as SOD, CAT, activity and glutathione level in almost all tissue of high fat induced rats compared with non-fat animals. On the other hand, non-enzymatic oxidative stress parameters lipid peroxidation marker MDA level increased in these tissues. Several studies have been shown that high fat diet induced decrease antioxidant capacity in different organs. In present study the significant decrease in MDA concentration and significantly increase in other antioxidant parameters shown in the standard group, administered with standard antioxidant agent Vit E (α -tocopherol acetate), indicates the integrity and validity of our experimental procedure by showing the expected protective effects of Vit E against all antioxidant parameters. In the present study showed that the 50% ethanolic extract of compound shown to exert its antioxidant effect mainly by ROS generation blockade due to increased SOD and catalase and Glutathion Reductase activity. However, Lipid peroxidation also plays an important role in its antioxidant activity.

CONCLUSION

The combination of the plant ethanolic extract exhibited significant antioxidant, activities as compared with alpha tocopherol acetate (Standard). This study could provide a new basis on the using of herbal medicinal plants in combination for the effective treatment by antioxidant mechanism.

Acknowledgements: The author is grateful to Department of Biochemistry, Faculty of Life Science AMU, Aligarh, for approving the study by Departmental Institutional Animal Ethics Committee (IAEC), also the author would like to thank Mr. Sehzad AMU, Aligarh, for helping in animal handling.

REFERENCES

1. Anonymous, The Unani Pharmacopia of India, Central council for research in Unani Medicine, Part. I. Vol. I. 2007, 50-51.
2. Evans, WC., Trease and Evans Pharmacognosy, Saunders Elsevier Edinburgh London New York Philadelphia St Louis Sydney Toronto, 16thedn. 2009, 188.
3. Nabi MG. Makhzan Mufradat wa Murakkabat-e-Azam. Ma'roof ba Khawasul Advia (Narain Das Jangali Mal).jayyed Barqi Press, Ballimaran, Delhi. 1958; 141.
4. Khan MA. Muheet-e-Azam.Matbaa Nizamee. Kanpur. Part. I. Vol. II, 1313, 83-84.
5. Ibn Sina, "Alqanoon Fit-Tib" (Urdu translation by Golam Husain Kantoori), Kocha Chilan Daryi Ganj New Delhi, Vol. II, 2007, 62-63.
6. Anwar F, Latif S, Ashraf M, Gilani AH. Moringa oleifera: a food plant with multiple medicinal uses. Phytotherapy Research. 21(1), 2007, 17-25.



7. Dymock W, Warden CJH, and Hooper D. Pharmacographia Indica. A History of the Principal Drugs. Pub. The Institute of Health and Tibbi Research, Hamdard National Foundation. Pakistan. Vol. I, 1972, 107-108. & 337-338.
8. Nadkarni KM. Indian Materia Medica. 3rd ed. Popular Book Depot Bombay 7, Dhootapapeshwar Prkaashan, Ltd Vol. 1, 1982, P. 811-816. & 1126-1129.
9. Khory RN and Katarak, NN. Materia Medica of India and Therapeutics, 3rd Reprint Edition. Pub. Neeraj Publishing House Delhi- 110052, 1993, 235-236.
10. Kirtikar KR and Basu, BD. Indian Medicinal Plants. Pub. International Book Distributors, Dehradun. Vol. III, 1995, 1858-1861.
11. Chopra IC, Handa KL, and Kapur LD. Indigenous Drugs of India. U. N. Dhur and Sons Private Limited, 15, Bankim Chatterjee Street Calcutta-12. 2nd Edition, 1958, 515.
12. Ghani MN. Khazayinul Advia. Central council for Research In Unani Medicine, Ministry of Health and Family Welfare, Govt. of India, New Delhi. 1st Edition. Vol. IV, 2010, 507-511.
13. Anonymous. The Unani Pharmacopoeia of India. Central council for Research, In Unani Medicine, Part. I. Vol. V, 2008, 88-89.
14. Hirose N, Inoue T, Nishihara K, Sugano M, Akimoto K, Shimizu S *et al.* Inhibition of cholesterol absorption and synthesis in rats by sesamin. J Lipid Res 32, 1991, 629-638.
15. Bedigian D, Harlan JR. Evidence for cultivation of sesame in the ancient world. Econ Bot 40, 1986, 137-154.
16. Shahidi F, Liyana-pathirana CM, Wall D. Antioxidant activity of white and black sesame seeds and their hull fractions. Food Chem 99, 2006, 478-483.
17. Liu Z, Saarinen NM, Thompson LU. Sesamin is one of the major precursors of mammalian lignans in sesame seed (*Sesamum indicum*) as observed *in vitro* and in rats. J Nutr 136, 2006, 906-912.
18. Ghafoorunissa, Hemalatha S, Rao MV. Sesame lignans enhance antioxidant activity of vitamin E in lipid peroxidation systems. Mol Cell Biochem 262, 2004, 195-202.
19. Costa FT, Neto SM, Bloch C, Franco OL. Susceptibility of human pathogenic bacteria to antimicrobial peptides from sesame kernels. Curr Microbiol 55, 2007, 162-166.
20. Yokota T, Matsuzaki Y, Koyama M, Hitomi T, Kawanaka M, Enoki-Konishi M. Sesamin, a lignan of sesame, downregulates cyclin D1 protein expression in human tumor cells. Cancer Sci 98, 2007, 1447-1453.
21. Visavadiya NP, Narasimhacharya AV. Sesame as a hypocholesteremic and antioxidant dietary component. Food Chem. Toxicol 46, 2008, 1889-1895.
22. Bhattacharjee, A.K., Das, A.K. Phytochemical survey of few Mysore Plants. Economic Botany, 23(3), 1969, 274-276.
23. Freirich EJ, Gehan EA, Rall DP, Schmidt LH, Skipper HE. Quantitative comparison of toxicity of anti ulceragents in mouse, rat, hamster, dog, monkey and man. Cancer chemotherapy Report 50(4), 1966, 219-244.
24. Kimuru, Y., Rhminani, H., Okuola, H. Effect of extract of Oyster on lipid metabolism in rats. J Ethnopharm. 1998, 117-123.
25. Friedewald WT., Levy RI., Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. Clinical Chemistry. 18(6), 1972, 499-502.
26. Malstrom, B., Andreasson, L., and Reinhammer, B. in the enzymes, Boyer, P., editor, XIIB, Academic Press, New York, 1975, 533.
27. Fridovich, I., Biological aspects of Superoxide radical and Superoxide dismutase. Biochemical and Medical Aspects of Active Oxygen. Asada K and Hayaish O, editor, Baltimore: University Park Press, 1977, 171-176.
28. Aebi, H.E. Catalase *in vivo*. Methods in Enzymology. L Packer, editor. N.York: Academic Press; Vol. 105, 1984, 121.
29. Yagi, K. Simple assay for the level of total lipid peroxidation in serum or plasma. Methods in Molecular Biology, 108, 1998, 101-6.
30. Armstrong, D., Browne, R., A systemic approach to laboratory technology, clinical correlation and antioxidant therapy. Free radicals in Diagnostic Medicine, 366, 1994, 43-58.
31. Spiteller, G. Lipid peroxidation in aging and age-dependent diseases, Exp, Geront. 36, 2001, 1425-1457
32. Evans, P., Halliwell, B. Free Radicals and hearing, causes, consequence and criteria. Ann. N.Y. Acad. Sci, 884, 1999, 1940.
33. Noeman, S.A., Hamooda, H.E., Biomedical study of oxidative stress markers in liver, kidney and heart of high fat diet induced obesity in rats, Diabetology & metabolic syndrome, 3, 2011, 1-17.

Source of Support: Nil, Conflict of Interest: None.

