Review Article

Nitrosamine Impurities in Processed Meats: Formation, Detection, and Risk Mitigation Strategies

R. Vijayageetha, Sangavi G*, C.N. Nalini, Manasa D, Bhagyalatasatapathy, Prem Kumar V, Anitha T.
Department of Pharmaceutical Analysis, C.L.Baid Metha College of Pharmacy, 2/305, Rajiv Gandhi Salai, OMR, Thoraipakkam, Chennai-6000097, Tamil Nadu, India.

***Corresponding author's E-mail:** sangavi292@gmail.com

Received: 05-05-2026; **Revised:** 22-07-2026; **Accepted:** 29-07-2026; **Published online:** 20-08-2026.

ABSTRACT

Nitrosamines are a group of N-nitroso compounds generally recognized as potent carcinogens. They may be formed during the processing, preservation, or even digestion of food, especially in processed meats containing added nitrites. Under acidic conditions, nitrites can combine with amines to form volatile and non-volatile nitrosamines such as NDMA, NDEA, and NPYR. These have been linked to an increased risk of malignant tumours of the liver, lungs, and digestive tract, as shown in many epidemiological and toxicological studies. Due to the development of appropriate analysis methods, including primarily GC-MS and LC-MS/MS techniques, the identification and quantification of a wide series of nitrosamines in meat products can now be performed with high sensitivity. Nitrite reduction, addition of antioxidants, or re-evaluation of cooking and/or processing procedures could decrease the exposure to nitrosamines. Moreover, ensuring a greater awareness by consumers and stricter regulatory measures is necessary for minimizing exposure and protecting public health against noxious substances like nitrosamines.

Keywords: Nitrosamines, chemistry, carcinogens, food preservation, food safety, risk assessment.

INTRODUCTION

International Agency for Research on Cancer (IARC) defines, "Nitrosamines, or N-nitroso compounds, are a group of chemical compounds that have a nitroso group (–N=O) attached to a nitrogen atom. Many N-nitrosamines are strong cancer-causing agents in experimental animals and are thought to increase the risk of cancer in humans. They can form in the body from nitrite and amines, or during food processing, preservation, and cooking—especially in cured meats" [1,2].

When secondary or tertiary amines react with nitrosating agents like nitrite/nitrogen oxides, they form N-nitrosamines. These are chemical molecules that have a nitroso group (-N=O) attached to amine. The production of nitrosamines, many of which are known to cause cancer, can occur from these agents. These agents often show up during the curing and processing of meats in the stomach due to the acidic environment. Volatile nitrosamines and non-volatile nitrosamines can form in processed meats when nitrite reacts with amines during processing, especially at higher temperatures, such as during frying/smoking. The body can produce endogenous nitrosamines through the nitrosation of nitrite and amines during digestion. Acidic conditions in the stomach can further promote this process. There is considerable evidence of nitrosamines in food, particularly in meat products that are often processed with nitrite. Nitrosamines are a group of compounds, and most of them are cancer-causing in nature [3].

Secondary and tertiary amines, amides, peptides, ureas, and guanidines are present in common foods, especially in protein-rich sources like meat, fish, and dairy, makes N-nitroso compounds important from a nutrition and public health standpoint. Although N-nitroso compounds are generally low in food and beverages, their precursors are commonly found in our diet. Because of this, N-nitroso compounds, including nitrosamines and nitrosamides, are recognized as one of the most significant classes of chemical carcinogens. Studies show that over 90% of tested N-nitroso compounds are carcinogenic in at least one animal species. Many have effects on multiple organs, and some are very potent even at low doses. Note that exposure to nitrosamines does not only come from eating pre-formed compounds. Humans can also be exposed through internal processes. Nitrite, which comes from dietary nitrate or food additives, can react with these amine precursors within the body, creating N-nitroso compounds, especially in acidic conditions of the stomach. The interest in nitrosamines began in the late 1950s and 1960s when researchers linked volatile nitrosamines, like NDMA, to cancer in rodents. Later research expanded this concern to non-volatile nitrosamines and ways the body can produce them, showing that normal digestion can create potentially carcinogenic N-nitroso compounds from regular meals. This interaction between precursors, nitrite/nitrate intake, and stomach chemistry has influenced both food safety regulations, such as limits on nitrite in cured meats, and nutrition advice, like encouraging diets rich in antioxidants to reduce nitrosation [4,5].

Historically, their detection in the 1970s in beer, processed meats, and tobacco smoke raised public health concerns and led to regulatory action. The widespread presence of precursors—amines from proteins and nitrite from food additives and vegetables—creates a constant potential for nitrosamine formation in daily life. By the early 1970s, researchers found N-nitrosodimethylamine (NDMA) in various foods & beverages, but its discovery in beer raised specific concerns. Beer is widely consumed, often daily, by many people. NDMA is a strong carcinogen in lab animals,

©Copyright protected. Unauthorised republication, reproduction, distribution, dissemination and copying of this document in whole or in part is strictly prohibited.

capable of causing tumors in the liver and other organs at very low doses. This raised worries that lifelong exposure from beer consumption could contribute to cancer risk in humans. During malting, in the beer production, the breakdown of barley proteins released dimethylamine (DMA).

The use of direct-fired kilns introduced nitrogen oxides (NOx) from combustion gases. When DMA and NOx came into contact in the warm, moist conditions of kilning, chemical nitrosation reactions took place, producing NDMA[6-8]. Members of the larger group of nitroso compounds, which includes nitrosamines, have been researched extensively since the late 1950s. Laboratory animals first showed signs of their strong ability to cause cancer during this time.

1. Pre-formed nitrosamines are found in foods, drinks, tobacco, industrial products, cosmetics, and even medicines.

2. The body can also create nitrosamines when amine-containing compounds (from dietary proteins or drugs) react with nitrosating agents in the stomach or other biological environments [8-10].

Some N-nitroso compounds are active even at micrograms per kilogram, making them some of the strongest environmental carcinogens known. Many of them can cause mutations, which means they can form DNA adducts and lead to gene mutations. They are especially harmful to certain organs, notably the liver, lungs, esophagus, and bladder.

The nitrosamines are not only a food safety issue but also a problem that spans multiple sectors, including environmental contamination, workplace exposure, and pharmaceutical production. It points out that while modern regulations have lowered their levels in many products, ongoing monitoring, improved testing methods, and preventive steps are necessary to keep public exposure as low as reasonably possible[10-12].

Nitrosamines are a significant group of environmental and dietary carcinogenic agents. They can be created artificially during food processing or produced naturally in the human body. Many are strong carcinogens in laboratory animals, causing tumors in several organs. Their ability to cause cancer depends on how they are transformed into DNA-reactive forms in the body. The seven nitrosamines that are often found in food are mentioned below [13-15],

1. NDMA expanded as (N-nitrosodimethylamine)
2. NDEA expanded as (N-nitrosodiethylamine)
3. NPYR expanded as (N-nitrosopyrrolidine)
4. NPIP expanded as (N-nitrosopiperidine)
5. NMOR expanded as (N-nitrosomorpholine)
6. NDPA expanded as (N-nitrosodi-n-propylamine)
7. NDBA expanded as (N-nitrosodi-n-butylamine)

Types of nitrosamine impurities

The IARC Classification of nitrosamines are discussed in Figure 1. And details in Table1.The nitrosamine types include volatile nitrosamines and non-volatile nitrosamines. The nitrosamine levels ranging depending on their type in nitrite-processed meat products.

Non-volatile nitrosamines are particularly found in high amounts. Volatile nitrosamines, such as N-nitrosodimethylamine (NDMA) & N-nitrosopyrrolidine (NPYR), are usually strong carcinogens. In contrast, non-volatile nitrosamines, like N-nitrososarcosine (NSAR), are weak carcinogens, while others, including N-nitrosohydroxyproline (NHPRO), N-nitrosoproline (NPRO), N-nitroso-2-methyl-thiazolidine-4-carboxylic acid (NMTCA) & N-nitroso-thiazolidine-4-carboxylic acid (NTCA), are thought to be non-carcinogenic. However, the idea that non-volatile nitrosamines such as NTCA, NPRO, NMTCA & NHPRO are not cancer-causing still to be confirmed through real toxicological studies in vivo.

Theoretically, nitroso compounds can be decarboxylated to their carcinogenic forms (NThZ, NPYR, NMThZ, and NHPYR) either through microbial activity in the large intestine or during heat treatment. Until it is checked whether these presumed non-carcinogenic nitrosamines lead to the negative effects in health linked to processed meat consumption, their formation should be minimized as much as possible.

Nitrosamines found in foods include NDMA (N-nitrosodimethylamine), NDEA (N-nitrosodiethylamine), NDBA (N-nitrosodibutylamine), NPIP (N-nitrosopiperidine), NPYR (N-nitrosopyrrolidine), NMOR (N-nitrosomorpholine), NDPhA (N-nitrosodiphenylamine), NPRO (N-nitrosoproline), and NSAR (N-nitrososarcosine). The International Agency for Research on Cancer (IARC), under the World Health Organization (WHO), classifies nitrosamines based on their ability to cause cancer in human beings [16-18].



Figure 1: IARC Classification of nitrosamines

©Copyright protected. Unauthorised republication, reproduction, distribution, dissemination and copying of this document in whole or in part is strictly prohibited.

Table 1: Nitrosamine impurities and their IARC classification

Nitrosamine	IARC Classification	Details
NDMA (N-Nitrosodimethylamine)	Group 2A	Probable human carcinogen; induces liver, lung, and kidney tumors in animals
NDEA (N-Nitrosodiethylamine)	Group 2A	Probable human carcinogen; DNA alkylation, liver carcinogenesis
NPYR (N-Nitrosopyrrolidine)	Group 2B	Possible human carcinogen; formed during frying of bacon
NPIP (N-Nitrosopiperidine)	Group 2B	Found in processed meat with pepper or spices
NMOR (N-Nitrosomorpholine)	Group 2B	Found in food and tobacco smoke; induces liver and kidney tumors
NDBA (N-Nitrosodibutylamine)	Group 2B	Occurs in rubber, cured meats
NSAR (N-Nitrososarcosine)	Group 3	Found in cured and processed meats

Chemistry of nitrosamines

The general structure of Nitrosamine is R_2N, NO, where R represents alkyl groups. The nitroso group, N=O, attaches to a secondary amine nitrogen. The flat structure around the nitroso group depends on the lengths of the N–N and N–O bonds. For instance, the N–N and N–O bond lengths for nitrosamines like NDMA are about 132 and 126 pm [19-21].

Micelle-Catalyzed Reaction

Cationic micelles, such as cetyltrimethylammonium chloride, can speed up the nitrosation reaction of hydrophobic secondary amines like dibutylamine. This reduces the effective pKa and brings reagents closer together at the interface, increasing nitrosamine production by roughly 100 times [22].

Neutral Or Basic Conditions Via Carbonyl Catalysts

Carbonyl molecules, such as formaldehyde, can create nitrosamines even at nearly neutral pH levels. These compounds generate iminium intermediates that nitrite then attacks to form nitrosamines [23].

Nitrosation Via Oxidation in Reactive Nitrogen Species

In the presence of free radicals, reactive species like peroxynitrite ($ONOO^-$) can nitrosate secondary amines, producing N-nitrosamines even in less acidic conditions [24].

Mechanism

$NO_2^- + H^+ \rightarrow HNO_2$

$2HNO_2 \rightleftharpoons N_2O_3 + H_2O$

$N_2O_3 \rightarrow NO^+ + NO2^-$

$R_2NH + NO^+ \rightarrow R_2N–NO + H^+$

Nitrous acid (HNO2) is developed by the reaction of nitrite with protons in acidic settings, like the stomach or during meat curing. When nitrous acid breaks down, it produces unstable nitrosating species. At this point, nitrous acid (HNO2) reaches equilibrium and generates dinitrogen trioxide (N2O3) and nitrosonium ions (NO+). The nitrosating agents, NO^+ or N_2O_3, are attacked by a secondary amine (R_2NH), which acts as a nucleophile. This process creates nitrosamines ($R_2N–NO$). The chemistry behind nitrosamine formation is often quite complex. The specific conditions of the reaction, including the amounts of reactants, catalysts, and inhibitors, as well as the presence of other processes that might compete for the nitrosating agents, determine the final products from the various reactions that can happen at the same time [25-29].

NITROSAMINES IN PROCESSED MEATS:

Environmentally, particularly in food and water, N-nitrosamines can develop whenever secondary, primary, or even tertiary amines come into contact with nitrite. N-nitrosamines may also be formed by ingesting and exposing a mixture of meals, water, medications, and medical preparations that include both reactants to the stomach's acidic environment. The concentration of nitrite used as additive, residual nitrite, ascorbate, ascorbic acid, or a-tocopherol, nitrosamine precursors, pre-processing conditions and procedures, the adipose tissue percentage, the catalysts present or inhibitors of nitrosation reaction, the process of smoking, and the method of culinary art all influence the formation of nitrosamines in meat products [30-34]. The nitrosamine formation and detection pathway is given in Figure 2.



Figure 2: Pathway of Nitrosamine formation and detection

©Copyright protected. Unauthorised republication, reproduction, distribution, dissemination and copying of this document in whole or in part is strictly prohibited.

In chemical and energy industries, amine-based solvents are widely used for gas scrubbing (e.g., removing CO_2 or NOx). These systems often operate at high temperatures and in oxygen-rich, NOx-containing environments — conditions favorable for oxidative nitrosation. Regulatory frameworks focus on preventing acidic nitrosation in foods and pharmaceuticals, but oxidative pathways are less controlled and potentially underrecognized. Volatile secondary amines released from industrial activities, agriculture, or natural sources can undergo oxidation in the atmosphere, especially in the presence of nitrogen oxides (NOx), leading to nitrosamine formation. In living systems, oxidative nitrosation can occur during inflammatory processes, where reactive nitrogen species (RNS) and reactive oxygen species (ROS) are both developed. Unlike acid-mediated nitrosation, oxidative nitrosation involves radical intermediates and metal-catalyzed oxidation steps, expanding the range of conditions under which nitrosamines can form [35-39].

Nitrosamines in processed foods may result from high-temperature cooking methods and contact with combustion gases, such as toasting, smoking, grilling or roasting. Many factors influence how much nitrosamine is produced during thermal food processing. These include temperature, how long the food is treated, the distance from the heating source, the content of fat, and type of fuel used. The nitrosamine formation in meats is a complex process, and many factors can affect the nitrosation reaction. The main sources of nitrosyl donors are nitrogen oxides from smoking and sodium nitrite from curing. N-nitrosamines form when these nitrosyl donors react with secondary and tertiary amines. There are several ways that secondary amines can get into food products. It is thought that biogenic amines and other substances produced during protein breakdown in meat are important sources of amine precursors. According to research, humans can tolerate exposure to the most volatile nitrosamines at levels of 5 to 10 µg kg^{-1} of body weight [40-43]. The various reasons for the formation of nitrosamine in Figure 3.



Figure 3: Formation of Nitrosamine

The levels of nitrosamine chemicals in raw meats can differ based on several factors, such as the animal species, their diets, and seasonal changes. Various complex chemical reactions occur during production, storage, packaging, and cooking, affecting how nitrosamines form in beef products. Some studies have connected processed beef consumption to an increased risk of stomach cancer.

Although only a few studies have directly looked at the relationship between nitrosamine consumption and the risk of stomach cancer, nitrosamines could be responsible for this connection. Nitrocompounds often appear in food at very low levels. This reduces human exposure to them. However, due to their strong potential to cause cancer, we should pay attention to these chemicals. Processed meats like sausages, bacon, ham often use nitrites for flavour and colour enhancement and to prevent spoilage and microbial growth. Cooking and smoking the meats at high temperature can increase the nitrosamine formation in the meats. Nitrosamines may be formed in the stomach due to their acidic environment. Factors like pH, cooking temperature, additives, storage time, microbial activity influencing the formation of nitrosamine [44-48].

Reasons for the synthesis of nitrosamines in different foods will differ. Processed meats, smoked foods, cured foods, dried and fermented foods, fried and grilled foods are different foods that contain nitrosamines. Processed meats include bacon, ham, sausages, hot dogs. The reaction between the nitrites (preservative) and the amines in those meats forms the nitrosamines. Formation of nitrosamine due to smoke chemicals containing nitrogen compounds in smoked foods. Smoked foods include smoked fish, smoked meats, and BBQ meats. Salami, pepperoni, cured beef are considered to be cured foods; the nitrites used in curing react with amines during storage and cooking, leading to the synthesis of nitrosamines. The bacterial activity in dried fish can increase the nitrosation reaction of amines and then nitrosamine synthesis. High-temperature cooking in grilled burgers induces nitrosation reactions, leading to nitrosamine formation [49-52].

Advances in detection and quantification of nitrosamines in processed meats

Analytical techniques for detecting, measuring, and comparing nitrosamines in different types of meat have improved significantly since the carcinogenic risk of these compounds in processed meats was recognized.

Nitrosamine levels in different products

Occurrence of nitrosamine impurities in meats is tabulated based on their type of nitrosamine impurity, including their detection methods [53-65] in Table 2. Figure 4. speaks on the levels of nitrosamine impurities in sausages.

©Copyright protected. Unauthorised republication, reproduction, distribution, dissemination and copying of this document in whole or in part is strictly prohibited.

Table 2: Occurrence of nitrosamine impurities in meat products and their detection methods

Name	Structure	Occurrence in Meats	Level (µg/kg)	Detection Methods
NDMA N-nitrosodimethylamine		Bacon, ham, hot dogs, sausages	0.3–2.1	GC-MS, LC-MS/MS
NDEA N-nitrosodiethylamine		Bacon, sausages, smoked meats	0.2–0.7	GC-MS, LC-MS/MS
NDPA N-nitrosodipropylamine		Cured meats, cheese, sausages	<0.1–0.4	GC-MS, LC-MS/MS
NDBA N-nitrosodibutylamine		Cured meats, smoked sausages	<0.1–0.3	GC-MS, LC-MS/MS
NPYR N-nitrosopyrrolidine		Bacon, sausages, cured meats	Up to 15.2	GC-MS
NPIP N-nitrosopiperidine		Sausages, smoked meats	Up to 1.7	GC-MS
NMOR N-nitrosomorpholine		Rare/cured meats	<0.1–0.5	GC-MS
NTCA N-nitrosothiazolidine-4-carboxylic acid		Smoked beef, sausage, corned beef	Up to 4227	LC-MS/MS
NMTCA N-nitroso-2-methyl-thiazolidine-4-carboxylic acid		Sausage, corned beef	20–989	LC-MS/MS



Figure 4: Levels of nitrosamine impurities in Sausage

Modern Detection Techniques

- *Liquid Chromatography tandem Mass Spectrometry:*

Extensively used for its sensitivity, selectivity, and ability to analyze multiple nitrosamine species simultaneously. Main tool for non-volatile, polar nitrosamines. Recent advances include QuEChERS-based sample preparation combined with LC-MS/MS, which allows rapid, accurate detection of up to 11 nitrosamine types in various processed meats and fish products. Extraction involves homogenizing samples with solvents like acetonitrile, centrifugation, filtration, and precise injection into the detection system (often using Electrospray ionization).

©Copyright protected. Unauthorised republication, reproduction, distribution, dissemination and copying of this document in whole or in part is strictly prohibited.

- *Gas Chromatography Mass Spectrometry:*

Effective for volatile nitrosamines (e.g., NDMA, NDEA). Extraction methods—such as ultrasonic treatment or autoclaving—improve recovery for accurate quantification. Solid Phase Microextraction (SPME) and headspace sampling enable efficient extraction and concentration of nitrosamines from meat matrices for direct injection are advanced sample preparation techniques. Coupled with selective ion monitoring, GC-MS offers low detection limits (~0.2 µg/kg), high specificity and simultaneous quantification of multiple compounds.

- *High Performance Liquid Chromatography:*

Used for simultaneous identification of volatile nitrosamines with direct UV detection, after autoclaving and organic solvent extraction. Capable of quantifying several nitrosamines in a single run. Validated for accuracy, linearity, and precision for food safety monitoring.

Table 3: Analytical techniques for nitrosamine determination

Method	Chromatographic parameters	Impurities	Retention time (min)	Reference

GC-MS	• **Sample Preparation:** Extraction with organic solvent like dichloromethane, sometimes clean-up with phosphate buffer.
	• **Column:** Non-polar capillary column (e.g., DB-5, 30 m × 0.25 mm, 0.25 µm film thickness).
	• **Injection:** Splitless, injector temp ~200–250°C.
	• **Oven Program:**
	  • Initial: 40–60°C (hold 2–5 min)
	  • Ramp 10°C/min to 220–250°C (hold 5–10 min)
	• **Carrier Gas:** Helium, 1 mL/min.
	• **Detection:** Chemical ionisation (methanol reagent) or electron impact.
	• **LOD/LOQ:** 0.15–0.37 µg/kg (LOD), 0.5–1.24 µg/kg (LOQ)
		NDMA, NMEA, NDEA, NPYR, NDPA, NMOR, NPIP, NDBA, NDPH.	NDMA - 4.11, NMEA - 5.29, NDEA - 6.40, NPYR - 9.22, NDPA - 9.25, NMOR - 9.40, NPIP - 10.25, NDBA - 14.91, NDPH - 19.75.	66

LC-MS/MS	• **Sample Preparation:** Extraction with acetonitrile, homogenisation, centrifugation, membrane filtration. • **Column:** C18 reversed-phase column (e.g., 250 mm × 4.6 mm, 5 µm). • **Mobile Phase:** Gradient of water (0.1% formic acid) and acetonitrile. • **Flow Rate:** 0.7–1.0 mL/min. • **Column Temp:** Ambient or controlled (30–40°C). • **Injection Volume:** 10 µL. • **Detection:** ESI (electrospray ionisation), MRM (multiple reaction monitoring). • **LOD:** Often sub-µg/kg	NTCA, NMTCA, NDEA.	NTCA - 2.65, NMTCA - 3.25, NDEA - 3.944.	67
RP-HPLC-UV	• **Column:** C18 • **Mobile Phase:** Gradient, 0%–90% solvent B (e.g., acetonitrile) over 10 min. • **Flow Rate**: 1.0 mL/min. • **Column Temperature:** 80°C • **Detection Wavelength:** 231 nm • **Sample Temperature:** 15°C • **Injection Volume:** 20 µL.	NDMA, NMEA, NDEA, NDPA, NDBA, NPIP, NPYR.	NDMA - 0.95, NMEA - 2.1, NDEA - 3.03, NDPA - 4.2, NDBA - 4.79, NPIP - 8.32, NPYR - 9.87.	68
LC-APCI-MS/MS	• **Column:** C18 150 × 4.6mm, 3.0µ column • **Mobile Phase:** Gradient, 0.1% formic acid in water as mobile phase A and 0.1% formic acid in methanol as mobile phase B • **Flow Rate:** 0.6 mL/min • **Run Time:** 25mins • **Column Temperature:** 40°C • **Injection Volume:** 30 µL. • Nebulizing Gas Flow 3.00 L/min • Interface Temperature 275 °C • DL Temperature: 200ºC • Heat Block temperature 200°C • Drying Gas Flow 3.00L/min	NDMA, NDEA and N-Nitroso varenicline	NDMA – 6.55, NDEA - 14.21, N-Nitroso varenicline - 17.94	69

©Copyright protected. Unauthorised republication, reproduction, distribution, dissemination and copying of this document in whole or in part is strictly prohibited.

Table 3 compares the analytical techniques for nitosamine determination. Modern techniques like GC-MS and LC-MS/MS offer quick, reliable nitrosamine detection and measurement in different processed meats. The largest concentrations of nitrosamines are present in smoked and cured meats, especially smoked beef and sausages, which often exceed recommended safety limits. Validated methods ensure precise risk assessment and support ongoing food safety monitoring and regulatory compliance[70-76].

RISK MANAGEMENT AND CONSUMER SAFETY:

Nitrites are often used as preservatives in meats. Reducing or eliminating the use of nitrites in meats will reduce the formation of nitrosamine impurities. On substituting nitrites as preservatives, natural preservatives like celery powder with antioxidants can be used. The lowest possible levels of nitrite can be added as the limits are set by the regulatory agencies. To inhibit nitrosamine formation the nitrosation reaction must be inhibited. The inhibitors, either ascorbic acid (vitamin C) / tocopherols (vitamin E), are added to meats. Polyphenols from natural sources like green tea and rosemary are also added. Processing methods of the meats were improved. Either avoid high heat cooking like frying or grilling or use indirect smoking or low temperature processing. Alternative preservation techniques for the meats are used. Irradiation can reduce the breakdown of nitrosamines. To reduce the formation of precursors, modified atmosphere packing (MAP) and vacuum packing can be done. To reduce the amine content in the raw meat feed, modifications to animals can be done. The diet of the animal must be altered [77-80].

Limiting the levels of nitrite and nitrate used in the process of curing is an important regulatory task. The USDA has established an upper limit of 10 µg/kg for total volatile nitrosamines in beef products. In China, the maximum allowed amount of NDMA in processed beef is 3 µg/kg. The European Union closely monitors the levels of nitrosamines in meat and medications, and it limits the use of leftover nitrite and nitrate. Regular testing for nitrosamines in processed meats, labelling requirements for products containing nitrate or nitrite, and guidelines for meat composition, temperature, and smoking methods are all part of this regulatory control [81-83].

Increasing public awareness and educating people is essential. Customers should understand the risks of consuming too much processed meat, safer cooking methods, and the nutritional benefits of antioxidants. Encourage consumers to choose meat products that are free from nitrates and nitrites or are naturally cured. Instead of grilling or frying meats, they should steam or boil them. Eating a diet rich in fruits and vegetables helps provide vitamin C and other nitrosation blockers. Make sure clients know the difference between nitrites and nitrates. Encourage manufacturers to use phrases like "no added nitrites" or "uncured" when relevant [84-88].

CONCLUSION

Nitrosamines are a major class of chemical contaminants that form during food processing, preservation, and digestion. Processed meats are a primary source of exposure to nitrosamines. These compounds develop due to interactions between nitrites and amines, especially in hot or acidic conditions. There is strong evidence of toxicity and tumor formation in animal studies. Many nitrosamines, including NDMA and NDEA, are classified as likely human carcinogens. The importance of cured and smoked meat products for public health has been underscored by widespread contamination found through sensitive analytical methods like LC-MS/MS and GC-MS.

Despite the serious health risks associated with nitrosamines in processed meats, we have made significant progress in understanding their chemistry, detection, and control. Today's sophisticated analytical techniques, such as GC-MS and LC-MS/MS, allow for precise monitoring of even small amounts. This helps food producers and regulators act quickly. Importantly, research and industry efforts have demonstrated that we can greatly reduce nitrosamine formation. Effective strategies include lowering nitrite levels, using antioxidants like ascorbic acid, switching to different preservatives, and improving cooking and smoking methods.

ACKNOWLEDGMENTS:

All authors have made substantial contributions to the work reported in the manuscript. Authors would like to thank all the individuals for their expertise and assistance throughout all aspects of our study and for their help in writing the manuscript.

Source of Support: The author(s) received no financial support for the research, authorship, and/or publication of this article

Conflict of Interest: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

REFERENCES

1. Bouvard V, Loomis D, Guyton KZ, Grosse Y, Ghissassi FE, Benbrahim-Tallaa L, et al. Carcinogenicity of consumption of red and processed meat. Lancet Oncol [Internet]. 2015;16(16):1599–600. https://www.iarc.fr/en/media-centre/pr/2015/pdfs/pr240_E.pdf
2. IARC. Some N-Nitroso Compounds. https://publications.iarc.who.int/Book-And-Report-Series/Iarc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Some-Em-N-Em--Nitroso-Compounds-1978
3. Management Board members, Executive Director, Operational Management. EFSA confirms safe levels for nitrites and nitrates added to food [Internet]. European Food Safety Authority. 2017 [cited 2026 May 15]. https://www.efsa.europa.eu/en/press/news/170615
4. Management Board members, Executive Director, Operational Management. Nitrosamines in food raise a health concern [Internet]. European Food Safety Authority. 2023 [cited 2026 May 15]. https://www.efsa.europa.eu/en/news/nitrosamines-food-raise-health-concern

©Copyright protected. Unauthorised republication, reproduction, distribution, dissemination and copying of this document in whole or in part is strictly prohibited.

5. Thomas R, Tennant RE, Oliveira AAF, Ponting DJ. What makes a potent nitrosamine? Statistical validation of expert-derived structure-activity relationships. Chem Res Toxicol [Internet]. 2022;35(11):1997–2013. https://doi.org/10.1021/acs.chemrestox.2c00199
6. Sebranek JG, Cassens RG. NITROSAMINES: A REVIEWa. J Milk Food Technol [Internet]. 1973;36(2):76–91. https://doi.org/10.4315/0022-2747-36.2.76
7. Challis BC. Nutrition and nitrosamine formation. Proc Nutr Soc [Internet]. 1985;44(1):95–100. https://doi.org/10.1079/pns19850015
8. Scanlan RA. Formation and occurrence of nitrosamines in food. Cancer Res. 1983;43(5 Suppl):2435s–40s. https://pubmed.ncbi.nlm.nih.gov/6831466/
9. Bond T, Templeton MR. Nitrosamine formation from the oxidation of secondary amines. Water Sci Technol Water Supply [Internet]. 2011;11(3):259–65. https://doi.org/10.2166/ws.2011.027
10. Hecht SS. Approaches to cancer prevention based on an understanding of N-nitrosamine carcinogenesis. Proc Soc Exp Biol Med [Internet]. 1997;216(2):181–91. https://doi.org/10.3181/00379727-216-44168
11. Li Y, Hecht SS. Metabolic activation and DNA interactions of carcinogenic N-nitrosamines to which humans are commonly exposed. Int J Mol Sci [Internet]. 2022;23(9):4559. https://doi.org/10.3390/ijms23094559
12. Herrmann SS, Duedahl-Olesen L, Christensen T, Olesen PT, Granby K. Dietary exposure to volatile and non-volatile N-nitrosamines from processed meat products in Denmark. Food Chem Toxicol [Internet]. 2015;80:137–43. https://doi.org/10.1016/j.fct.2015.03.008
13. Xie Y, Geng Y, Yao J, Ji J, Chen F, Xiao J, et al. N-nitrosamines in processed meats: Exposure, formation and mitigation strategies. J Agric Food Res [Internet]. 2023;13(100645):100645. https://doi.org/10.1016/j.jafr.2023.100645
14. Metzler M. Nitrosamines. Toxicology and Microbiology. (Ellis Horwood Series in Food Science and Technology). Herausgegeben vonM. J. Hill. VCH Verlagsgesellschaft, Weinheim/VCH Publisher, New York/Ellis Horwood, Chichester 1988. 169 S., geb., DM 180.00. – ISBN 3-527-26708-5/0-89573-605-0; ISSN 0930-3332. Angew Chem Weinheim Bergstr Ger [Internet]. 1989;101(10):1434–1434. https://doi.org/10.1002/ange.19891011038
15. Park J-E, Seo J-E, Lee J-Y, Kwon H. Distribution of seven N-nitrosamines in food. Toxicol Res [Internet]. 2015;31(3):279–88. https://doi.org/10.5487/TR.2015.31.3.279
16. Deveci G, Tek NA. N-Nitrosamines: a potential hazard in processed meat products. J Sci Food Agric [Internet]. 2024;104(5):2551–60. https://doi.org/10.1002/jsfa.13102
17. Rot T, Kovačević D, Habschied K, Mastanjević K. N-nitrosamines in meat products: Formation, detection and regulatory challenges. Processes (Basel) [Internet]. 2025;13(5):1555. https://doi.org/10.3390/pr13051555
18. Lijinsky W. Chemistry and biology of N-nitroso-compounds. In: Coombs MM, Ashby J, Ricks M, editors. Chemistry and Biology of N-nitrosocompounds. Cambridge: Cambridge University Press; 1992.
19. Gray JI, Randall CJ. The nitrite/N-nitrosamine problem in meats: An update. J Food Prot [Internet]. 1979;42(2):168–79. https://doi.org/10.4315/0362-028X-42.2.168
20. Özbay S. Nitrosamine formation in processed meat products: A comprehensive analysis of ham varieties, including turkey and chicken, in the Turkish market [Internet]. 2023. https://doi.org/10.2139/ssrn.4663063
21. Herrmann SS, Granby K, Duedahl-Olesen L. Formation and mitigation of N-nitrosamines in nitrite preserved cooked sausages. Food Chem [Internet]. 2015;174:516–26. https://doi.org/10.1016/j.foodchem.2014.11.101
22. Breider F, Salihu I, von Gunten U. Formation of N-nitrosamines by micelle-catalysed nitrosation of aliphatic secondary amines. Environ Sci Process Impacts [Internet]. 2018;20(10):1479–87. https://doi.org/10.1039/c8em00335a
23. Pan Y, Breider F, Barrios B, Minakata D, Deng H, von Gunten U. Role of carbonyl compounds for N-nitrosamine formation during nitrosation: Kinetics and mechanisms. Environ Sci Technol [Internet]. 2024;58(10):4792–801. https://doi.org/10.1021/acs.est.3c07461
24. Masuda M, Mower HF, Pignatelli B, Celan I, Friesen MD, Nishino H, et al. Formation of N-nitrosamines and N-nitramines by the reaction of secondary amines with peroxynitrite and other reactive nitrogen species: comparison with nitrotyrosine formation. Chem Res Toxicol [Internet]. 2000;13(4):301–8. https://doi.org/10.1021/tx990120o
25. Mirvish SS. Formation of N-nitroso compounds: chemistry, kinetics, and in vivo occurrence. Toxicol Appl Pharmacol [Internet]. 1975;31(3):325–51. https://doi.org/10.1016/0041-008x(75)90255-0
26. Tricker AR, Preussmann R. Carcinogenic N-nitrosamines in the diet: occurrence, formation, mechanisms and carcinogenic potential. Mutat Res [Internet]. 1991;259(3–4):277–89. https://doi.org/10.1016/0165-1218(91)90123-4
27. Farvid MS, Sidahmed E, Spence ND, Mante Angua K, Rosner BA, Barnett JB. Consumption of red meat and processed meat and cancer incidence: a systematic review and meta-analysis of prospective studies. Eur J Epidemiol [Internet]. 2021;36(9):937–51. https://doi.org/10.1007/s10654-021-00741-9
28. Wainright T. The chemistry of nitrosamine formation: relevance to malting and brewing. J Inst Brew [Internet]. 1986;92(1):49–64. https://doi.org/10.1002/j.2050-0416.1986.tb04373.x
29. Iko Afé OH, Kpoclou YE, Douny C, Anihouvi VB, Igout A, Mahillon J, et al. Chemical hazards in smoked meat and fish. Food Sci Nutr [Internet]. 2021;9(12):6903–22. https://doi.org/10.1002/fsn3.2633
30. Larsson SC, Bergkvist L, Wolk A. Processed meat consumption, dietary nitrosamines and stomach cancer risk in a cohort of Swedish women. Int J Cancer [Internet]. 2006;119(4):915–9. https://doi.org/10.1002/ijc.21925
31. Colă F, Colă M. Researches on formation of nitrosamines in meat products. Annals of the University of Craiova - Agriculture. 2018;48(1):88–92.
32. Gilani PS, Fesahat M, Shariatifar N. Nitrosamine in meat and meat products: a review. J Food Saf Hyg [Internet]. 2024. https://doi.org/10.18502/jfsh.v9i4.14998
33. Kaban G, Polat Z, Sallan S, Kaya M. The occurrence of volatile N-nitrosamines in heat-treated sucuk in relation to pH, aw and residual nitrite. J Food Sci Technol [Internet]. 2022;59(5):1748–55. https://doi.org/10.1007/s13197-021-05186-2
34. Niklas AA, Herrmann SS, Pedersen M, Jakobsen M, Duedahl-Olesen L. The occurrence of volatile and non-volatile N-nitrosamines in cured meat products from the Danish market. Food Chem [Internet]. 2022;378(132046):132046. https://doi.org/10.1016/j.foodchem.2022.132046
35. Rywotycki R. The effect of baking of various kinds of raw meat from different animal species and meat with functional additives on nitrosamine contamination level. Food Chem [Internet]. 2007;101(2):540–8. https://doi.org/10.1016/j.foodchem.2006.02.012
36. Sen NP, Seaman SW, Burgess C, Baddoo PA, Weber D. Investigation on the possible formation of N-nitroso-N-methylurea by nitrosation of creatinine in model systems and in cured meats at gastric pH. J Agric Food Chem [Internet]. 2000;48(10):5088–96. https://doi.org/10.1021/jf990918d
37. Sallan S, Kaban G, Kaya M. Nitrosamines in sucuk: Effects of black pepper, sodium ascorbate and cooking level. Food Chem [Internet]. 2019;288:341–6. https://doi.org/10.1016/j.foodchem.2019.02.129

©Copyright protected. Unauthorised republication, reproduction, distribution, dissemination and copying of this document in whole or in part is strictly prohibited.

38. Li L, Wang P, Xu X, Zhou G. Influence of various cooking methods on the concentrations of volatile N-nitrosamines and biogenic amines in dry-cured sausages. J Food Sci [Internet]. 2012;77(5):C560-5. https://doi.org/10.1111/j.1750-3841.2012.02667.x

39. Souliotis VL, Henneman JR, Reed CD, Chhabra SK, Diwan BA, Anderson LM, et al. DNA adducts and liver DNA replication in rats during chronic exposure to N-nitrosodimethylamine (NDMA) and their relationships to the dose-dependence of NDMA hepatocarcinogenesis. Mutat Res [Internet]. 2002;500(1–2):75–87. https://doi.org/10.1016/s0027-5107(01)00301-3

40. De Mey E, De Klerck K, De Maere H, Dewulf L, Derdelinckx G, Peeters M-C, et al. The occurrence of N-nitrosamines, residual nitrite and biogenic amines in commercial dry fermented sausages and evaluation of their occasional relation. Meat Sci [Internet]. 2014;96(2 Pt A):821–8. https://doi.org/10.1016/j.meatsci.2013.09.010

41. Reinik M. Nitrates, nitrites, N-nitrosamines and polycyclic aromatic hydrocarbons in food: analytical methods, occurrence and dietary intake. 2007.

42. Zhang Y, Zhang Y, Jia J, Peng H, Qian Q, Pan Z, et al. Nitrite and nitrate in meat processing: Functions and alternatives. Curr Res Food Sci [Internet]. 2023;6(100470):100470. https://doi.org/10.1016/j.crfs.2023.100470

43. Lu J, Li M, Shen M, Xie J, Xie M. Advanced glycation end products and nitrosamines in sausages influenced by processing parameters, food additives and fat during thermal processing. Foods [Internet]. 2023;12(2):394. https://doi.org/10.3390/foods12020394

44. Herrmann SS. N-nitrosamines in processed meat products - analysis, occurrence, formation, mitigation and exposure. 2014. https://www.semanticscholar.org/paper/N-nitrosamines-in-processed-meat-products-%E2%80%93-and-Herrmann/35d64024f44d2a9bd283c38dfa5e70775b0b2ffa

45. Herrmann SS, Duedahl-Olesen L, Granby K. Occurrence of volatile and non-volatile N-nitrosamines in processed meat products and the role of heat treatment. Food Control [Internet]. 2015;48:163–9. https://doi.org/10.1016/j.foodcont.2014.05.030

46. Özbay S, Şireli UT. Volatile N-nitrosamines in processed meat products and salami from Turkey. Food Addit Contam Part B Surveill [Internet]. 2021;14(2):110–4. https://doi.org/10.1080/19393210.2021.1885502

47. Ahn H-J, Kim J-H, Jo C, Lee J-W, Yook H-S, Byun M-W. Effects of gamma irradiation on residual nitrite, residual ascorbate, color, and N-nitrosamines of cooked sausage during storage. Food Control [Internet]. 2004;15(3):197–203. https://doi.org/10.1016/s0956-7135(03)00047-1

48. Mavelle T, Bouchikhi B, Debry G. The occurrence of volatile N-nitrosamines in French foodstuffs. Food Chem [Internet]. 1991;42(3):321–38. https://doi.org/10.1016/0308-8146(91)90073-w

49. Niklas AA, Borge GIA, Rødbotten R, Berget I, Müller MHB, Herrmann SS, et al. Levels of nitrate, nitrite and nitrosamines in model sausages during heat treatment and in vitro digestion - The impact of adding nitrite and spinach (Spinacia oleracea L.). Food Res Int [Internet]. 2023;166(112595):112595. https://doi.org/10.1016/j.foodres.2023.112595

50. De Mey E, De Maere H, Paelinck H, Fraeye I. Volatile N-nitrosamines in meat products: Potential precursors, influence of processing, and mitigation strategies. Crit Rev Food Sci Nutr [Internet]. 2017;57(13):2909–23. https://doi.org/10.1080/10408398.2015.1078769

51. Nabizadeh S, Aeini K, Barzegar F, Arabameri M, Hosseini H, Kamankesh M, et al. Volatile N-nitrosamines in processed meat products: An approach for monitoring dietary exposure, assessing human risk, and evaluating variable correlations by principal component analysis and heat map. Food Chem Toxicol [Internet]. 2024;188(114649):114649. https://doi.org/10.1016/j.fct.2024.114649

52. Sheng S, Silva EM, Tarté R, Claus JR. Residual nitrite and nitrate in processed meats and meat analogues in the United States. Sci Rep [Internet]. 2025;15(1):3269. https://doi.org/10.1038/s41598-025-87563-x

53. Méndez Pérez D, González Alatorre G, Botello Álvarez E, Escamilla Silva E, Javier Alvarado JF. Solid-phase microextraction of N-nitrosodimethylamine in beer. Food Chem [Internet]. 2008;107(3):1348–52. https://doi.org/10.1016/j.foodchem.2007.09.064

54. Grebel JE, Mel Suffet IH. Nitrogen-phosphorus detection and nitrogen chemiluminescence detection of volatile nitrosamines in water matrices: optimization and performance comparison. J Chromatogr A [Internet]. 2007;1175(1):141–4. https://doi.org/10.1016/j.chroma.2007.09.073

55. Huang M-C, Chen H-C, Fu S-C, Ding W-H. Determination of volatile N-nitrosamines in meat products by microwave-assisted extraction coupled with dispersive micro solid-phase extraction and gas chromatography--chemical ionisation mass spectrometry. Food Chem [Internet]. 2013;138(1):227–33. https://doi.org/10.1016/j.foodchem.2012.09.119

56. Kocak D, Ozel MZ, Gogus F, Hamilton JF, Lewis AC. Determination of volatile nitrosamines in grilled lamb and vegetables using comprehensive gas chromatography - nitrogen chemiluminescence detection. Food Chem [Internet]. 2012;135(4):2215–20. https://doi.org/10.1016/j.foodchem.2012.07.002

57. Ozel MZ, Gogus F, Yagci S, Hamilton JF, Lewis AC. Determination of volatile nitrosamines in various meat products using comprehensive gas chromatography-nitrogen chemiluminescence detection. Food Chem Toxicol [Internet]. 2010;48(11):3268–73. https://doi.org/10.1016/j.fct.2010.08.036

58. Sonawane TS, Painginkar T, Bavadia R, Vasantharaju SG, Sathyanarayana MB, Ravi G. Navigating nitrosamines: Origin, detecting, analysing and regulating impurities in pharmaceuticals. Int J Appl Pharm [Internet]. 2024;1–11. https://doi.org/10.22159/ijap.2024v16i5.51376

59. Al-Kaseem M, Al-Assaf Z, Karabeet F. Development and validation of GC-FID method for the determination of volatile N-nitrosamines in meat. International Journal of Pharmaceutical Sciences Review and Research. 2014;25:59–64.

60. Al-Kaseem M, Al-Assaf Z, Karabeet F. Determination of seven volatile N-nitrosamines in fast food. Pharmacol Pharm [Internet]. 2014;05(02):195–203. https://doi.org/10.4236/pp.2014.52026

61. Crews C. The determination of N-nitrosamines in food: N-nitrosamines in food. Qual Assur Saf Crops Foods [Internet]. 2010;2(1):2–12. https://doi.org/10.1111/j.1757-837x.2010.00049.x

62. Roasa J, Liu H, Shao S. An optimised HS-SPME-GC-MS method for the detection of volatile nitrosamines in meat samples. Food Addit Contam Part A Chem Anal Control Expo Risk Assess [Internet]. 2019;36(3):396–404. https://doi.org/10.1080/19440049.2019.1571287

63. Lee HJ, Jung YS, Seo D, Kim E, Yoo M. Development and validation of QuEChERS-based LC-MS/MS method for simultaneous quantification of eleven N-nitrosamines in processed fish meat, processed meat, and salted fish products. Food Chem [Internet]. 2024;459(140281):140281. https://doi.org/10.1016/j.foodchem.2024.140281

64. Lu S, Wu D, Li G, Lv Z, Gong P, Xia L, et al. Facile and sensitive determination of N-nitrosamines in food samples by high-performance liquid chromatography via combining fluorescent labeling with dispersive liquid-liquid microextraction. Food Chem [Internet]. 2017;234:408–15. https://doi.org/10.1016/j.foodchem.2017.05.032

©Copyright protected. Unauthorised republication, reproduction, distribution, dissemination and copying of this document in whole or in part is strictly prohibited.

65. Abdullah ATM, Khan TA, Sharif M, Mazumdar RM, Rahman MM. Determination of dietary exposure and extraction efficiency of nitrosamine from cooked meat. Curr Res Food Sci [Internet]. 2022;5:491–7. https://doi.org/10.1016/j.crfs.2022.02.010

66. Scheeren MB, Sabik H, Gariépy C, Terra NN, Arul J. Determination of N-nitrosamines in processed meats by liquid extraction combined with gas chromatography-methanol chemical ionisation/mass spectrometry. Food Addit Contam Part A Chem Anal Control Expo Risk Assess [Internet]. 2015;32(9):1436–47. https://doi.org/10.1080/19440049.2015.1066037

67. Cintya H, Silalahi J, Putra EDL, Siburian R. Analysis of nitrosamines in processed meat products in Medan City by liquid chromatography-mass spectrometry. Open Access Maced J Med Sci [Internet]. 2019;7(8):1382–7. https://doi.org/10.3889/oamjms.2019.261

68. Al-Kaseem M, Al-Assaf Z, Karabeet F. A rapid, validated RP-HPLC method for the determination of seven volatile N-nitrosamines in meat. Pharmacol Pharm [Internet]. 2014;05(03):298–308. https://doi.org/10.4236/pp.2014.53037

69. Vidyamani U, Ramachandran D, Durga Babu M, Murthy Raju SS, Uma Maheswar K. Development and Validation of a Rapid and Sensitive LC-APCI-MS/MS Method for the determination of three N-nitrosamine impurities in Varenicline tartrate drug substance and drug products. Res J Pharm Technol [Internet]. 2024;1703–10. Available from: http://dx.doi.org/10.52711/0974-360x.2024.00270

70. Yurchenko S, Molder U. The occurrence of volatile N-nitrosamines in Estonian meat products. Food Chem [Internet]. 2007;100(4):1713–21. https://doi.org/10.1016/j.foodchem.2005.10.017

71. Manchuri KM, Shaik MA, Gopireddy VSR, Naziya Sultana, Gogineni S. Analytical methodologies to detect N-nitrosamine impurities in active pharmaceutical ingredients, drug products and other matrices. Chem Res Toxicol [Internet]. 2024;37(9):1456–83. https://doi.org/10.1021/acs.chemrestox.4c00234

72. Sun C, Wang R, Wang T, Li Q. Primary evaluation of nine volatile N-nitrosamines in raw red meat from Tianjin, China, by HS-SPME-GC-MS. Food Chem [Internet]. 2020;310(125945):125945. https://doi.org/10.1016/j.foodchem.2019.125945

73. Son W-Y, Kang M-H, Hwang J, Kim J-H, Dixit Y, Kim H-W. Hyperspectral imaging combined with machine learning can be used for rapid and non-destructive monitoring of residual nitrite in emulsified pork sausages. Foods [Internet]. 2024;13(19):3173. https://doi.org/10.3390/foods13193173

74. Shakil MH, Trisha AT, Rahman M, Talukdar S, Kobun R, Huda N, et al. Nitrites in cured meats, health risk issues, alternatives to nitrites: A review. Foods [Internet]. 2022;11(21):3355. https://doi.org/10.3390/foods11213355

75. Campillo N, Viñas P, Martínez-Castillo N, Hernández-Córdoba M. Determination of volatile nitrosamines in meat products by microwave-assisted extraction and dispersive liquid-liquid microextraction coupled to gas chromatography-mass spectrometry. J Chromatogr A [Internet]. 2011;1218(14):1815–21. https://doi.org/10.1016/j.chroma.2011.02.010

76. Chen S, Zhang Y, Zhao Q, Liu Y, Wang Y. Simultaneous determination for nine kinds of N-nitrosamines compounds in groundwater by ultra-high-performance liquid chromatography coupled with triple quadrupole mass spectrometry. Int J Environ Res Public Health [Internet]. 2022;19(24):16680. https://doi.org/10.3390/ijerph192416680

77. Sindelar JJ, Milkowski AL. Human safety controversies surrounding nitrate and nitrite in the diet. Nitric Oxide [Internet]. 2012;26(4):259–66. https://doi.org/10.1016/j.niox.2012.03.011

78. Honikel K-O. The use and control of nitrate and nitrite for the processing of meat products. Meat Sci [Internet]. 2008;78(1–2):68–76. https://doi.org/10.1016/j.meatsci.2007.05.030

79. Sen NP, Baddoo PA. Trends in the levels of residual nitrite in Canadian cured meat products over the past 25 years. J Agric Food Chem [Internet]. 1997;45(12):4714–8. https://doi.org/10.1021/jf9706641

80. Fashan WEI, Xinglian XU, Guanghong ZHOU. Determination method of volatile N-nioso compounds. Food Sci. 2008;29(7):479–83. https://www.spkx.net.cn/EN/Y2008/V29/I7/479

81. EFSA Panel on Plant Protection Products and their Residues (PPR), Ockleford C, Adriaanse P, Berny P, Brock T, Duquesne S, et al. Scientific Opinion addressing the state of the science on risk assessment of plant protection products for in-soil organisms. EFSA J [Internet]. 2017;15(2):e04690. https://doi.org/10.2903/j.efsa.2017.4690

82. Food US, Administration D, editors. Control of Nitrosamine Impurities in Human Drugs: Guidance for Industry. 2021. https://www.fda.gov/media/141720/download

83. EFSA Panel on Contaminants in the Food Chain (EFSA CONTAM Panel), Schrenk D, Bignami M, Bodin L, Chipman JK, Del Mazo J, et al. Risk assessment of N-nitrosamines in food. EFSA J [Internet]. 2023;21(3):e07884. https://doi.org/10.2903/j.efsa.2023.7884

84. Walters CL, Downes MJ, Edwards MW, Smith PL. Determination of a non-volatile N-nitrosamine on a food matrix. Analyst [Internet]. 1978;103(1232):1127–33. https://doi.org/10.1039/AN9780301127

85. Wogan GN, Tannenbaum SR. Environmental N-nitroso compounds: implications for public health. Toxicol Appl Pharmacol [Internet]. 1975;31(3):375–83. https://doi.org/10.1016/0041-008x(75)90259-8

86. Moradi S, Shariatifar N, Akbari-Adergani B, Molaee Aghaee E, Arbameri M. Analysis and health risk assessment of nitrosamines in meat products collected from markets, Iran: with the approach of chemometric. J Environ Health Sci Eng [Internet]. 2021;19(2):1361–71. https://doi.org/10.1007/s40201-021-00692-z

87. Tannenbaum SR. Preventive action of vitamin C on nitrosamine formation. Int J Vitam Nutr Res Suppl. 1989;30:109–13. https://pubmed.ncbi.nlm.nih.gov/2507690/

88. Tayel DI, Farrag NK, Aborhyem SM. Dietary intake and risk assessment of nitrosamine in processed meat products among medical staff during their night shift. Sci Rep [Internet]. 2025;15(1):1898. https://doi.org/10.1038/s41598-024-84059-y

For any questions related to this article, please reach us at: globalresearchonline@rediffmail.com

New manuscripts for publication can be submitted at: submit@globalresearchonline.net and submit_ijpsrr@rediffmail.com

©Copyright protected. Unauthorised republication, reproduction, distribution, dissemination and copying of this document in whole or in part is strictly prohibited.