



## EQUILIBRIUM STUDIES ON MIXED LIGAND COMPLEXES OF DRUG INDAPAMIDE WITH CHROMIUM AND COBALT METAL IONS

Pragati M. Deore<sup>1\*</sup>, Arun R. Khalkar<sup>2</sup>, B.R. Arbad<sup>3</sup>

1\*, 3. Department of Chemistry, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad, M.S, India.

2. Late G.N. Sapkal College of Engineering, Sapkal Knowledge Hub, Nashik, M.S, India.

\*Corresponding author's E-mail: pmdeore@gmail.com

Accepted on: 23-10-2011; Finalized on: 25-02-2012.


### ABSTRACT

Equilibrium studies on metal-ligand complex equilibria involving Chromium and Cobalt metal ions with drug, Indapamide with Amino acids Glycine and Glutamic acid in 80% (v/v) ethanol-water mixture at 30°C ± 0.1°C and ionic strength of 0.1M (NaClO<sub>4</sub>) has been studied. Formation of complex species with respect to pH have been discussed by Irving-Rossotti technique and evaluated by SCOGS computer program.

**Keywords:** Equilibrium constant, Ionic strength, pH, ΔlogK, SCOGS.

### INTRODUCTION

Drugs have various functional groups present in its structure, which can bind to metal ions present in human body<sup>1</sup>. Metal complexes of drugs are found to be more potent than parent drugs<sup>2</sup>. Chemistry of drugs attracts many researchers because of its applications in medicinal study. Interesting results have been reported earlier on complex formation reactions of drug-amino acid-metal ion mixed ligand complexes<sup>3-7</sup>. Expecting some useful information on mixed ligand complexes a detailed pH metric study involving drug Indapamide with Chromium and Cobalt metal ions has been carried out and discussed with results.



Indapamide

### MATERIALS AND METHODS

All the reagents used were of A.R. grade and all the solutions were prepared in 80% (v/v) ethanol-water mixture and standardized by known procedures<sup>8</sup>. Titrations were carried out using a digital pH meter (Elico model LI-127) in conjunction with combined electrode. All titrations were carried out at 30°C ± 0.1°C temp. All sets of solutions were titrated against 0.4N sodium hydroxide solution. The titration curves were plotted by using the experimental data. On the basis of these plots, proton-ligand and metal-ligand formation constants were calculated. Concentrations of total metal, total ligands, free metal, free ligands and various possible species that are formed during the complexation are calculated by using SCOGS computer program<sup>9,10</sup>.

### RESULTS AND DISCUSSION

#### Binary Complexes

Indapamide<sup>11-16</sup> is a mild diuretic and antihypertensive agent. It is the lipid soluble moiety, which distinguishes the activity of Indapamide from other diuretics. It has been shown to be a potent long acting antihypertensive agent when used along- with other therapeutic agents. The accepted chemical name for Indapamide is 3-(Aminosulfonyl)-4-chloro-N-(2, 3-dihydro-2-methyl-1H-indol-1-yl) benzamide.

The protonation constants  $K_1^H$  and  $K_2^H$  of drug Indapamide, amino acids Glycine and Glutamic acid and their metal-ligand formation constants ( $K_{ML}^H$  and  $K_{ML2}^{ML}$ ) have been determined (Table 1).

**Table 1:** Proton-ligand and metal-ligand stability constants in binary system

Ligands	$K_1^H$	$K_2^H$	Cr <sup>III</sup> (M <sub>1</sub> )		Co <sup>II</sup> (M <sub>2</sub> )	
			log K <sub>1</sub>	log K <sub>2</sub>	log K <sub>1</sub>	log K <sub>2</sub>
Indapamide (L)	3.23	9.97	7.28	3.49	7.09	3.59
Glycine (R <sub>1</sub> )	2.78	9.89	9.95	3.91	8.85	4.91
Glutamic acid (R <sub>2</sub> )	4.75	9.82	12.52	--	9.18	7.38

Indapamide drug molecule contains three different nitrogens. The calculated pK 3.23 can be assigned to amino nitrogen which readily undergoes the protonation in presence of acidic condition and subsequently deprotonates. The pK 9.97 can be assigned to amido group which undergoes self-deprotonation since it is connected to carbonyl group on one side and protonated nitrogen on other side, which stabilizes the corresponding nitrogen anion produced due to self-deprotonation.

Alongwith  $K_1^H$  and  $K_2^H$  values, also the stepwise metal-ligand formation constants of the ligands were determined for comparison with ternary systems. The complexing tendency of Cr<sup>III</sup> metal is found to be more than Co<sup>II</sup>. It shows that Cr<sup>III</sup> forms more stable five



membered chelate ring<sup>17</sup> with drugs used than other metal ions, it is due to higher charge on metal ion and smaller size<sup>18,19</sup>.

### Mixed ligand complexes

Complexes in which metal ion has two or more types of ligands in its coordinating sphere are called as mixed ligand complexes. The study of ternary complexes in solution provides simpler models for more complicated biochemical reactions<sup>20-23</sup>. Only 1:1:1 ternary complex have been used in this study to ensure the exclusive formation of the simplest ternary complex MLR. By considering the proton-ligand and metal-ligand constants of ligands, the species that exist in complexation equilibria have been plotted as a function of pH.




Figure 1: Species distribution curve for  $M_1LR_2$  system

It can be observed from Fig. 1 that mixed ligand curve of  $Cr(III)LR_2$  increases in the pH range of 4.5 to 5.9 with increase of pH. Whereas, the concentration for the formation of  $C_1$  and  $C_3$  shows continuous decrease with increasing pH which indicate the formation of  $Cr(III)L_4R_1$  complex and represented by  $C_8$ . Moreover, the maximum percentage of the formation of ternary complex is less than that of  $Cr(III)R_2$  binary complex and more than  $Cr(III)L$  binary complex, this indicates that the ternary complex is less stable as compared to  $Cr(III)R_2$  binary complex and more stable than  $Cr(III)L$  binary complex.




Figure 2: Percentage distribution curve for  $M_1LR_2$  system

Fig.2 shows that percentage distribution curve of free metal decreases with increase of pH, which indicates the involvement of metal ion in the complex formation process. Percentage concentration of free ligands  $FL_1$  and

$FL_2$  increases with increase of pH and this increase may be due to dissociation of ligand present in the system.

### CONCLUSION

The relative stabilities of the binary and ternary complexes are quantitatively expressed in terms of  $\beta_{111}, \beta_{20}, \beta_{02}, K_L, K_R, K_f$  and  $\Delta \log K$  values which are presented in Table No.2. The comparison of  $\beta_{111}$  with  $\beta_{20}$  and  $\beta_{02}$  of this system shows preferential formation of ternary complexes over binary complex of primary as well as secondary ligand. The considerably low positive value of  $K_L$  and  $K_R$  indicate less stability of ternary complex with respect to that of primary and secondary ligands. The  $K_f$  value of this complex is positive but less which indicate lower stability of ternary complex. The  $\Delta \log K$  value of this system indicates less stability of ternary complex.

Table 2: Stability constants of ternary complexes of Drug Indapamide

Metal ion	Amino Acid	$\beta_{111}$	$\beta_{20}$	$\beta_{02}$	$K_L$	$K_R$	$K_f$	$\Delta \log K$
Cr(III)	Glycine	14.48	10.77	13.86	7.20	4.53	1.17	-2.75
	Glutamic acid	15.81	10.77	12.52	8.52	3.28	1.36	-4.01
Co(II)	Glycine	13.14	10.68	13.77	6.04	4.27	1.07	-2.81
	Glutamic acid	17.13	10.68	16.56	10.04	7.94	1.25	0.86

### REFERENCES

- Walter S. Kittl, Bernd M. Rode, Complex formation of aliphatic dipeptides with zinc(II) and manganese(II), *Inorganica Chimica Acta*, 63,1982,47-52.
- Lancelot G., Hydrogen bonding of amino acid side chains to nucleic acidbases, *Biochimie*, Volume 59, Issue 7, 1977, 587-596.
- Dogan A., EsmaKilic, Potentiometric studies on the stability constants of some  $\alpha$ -amino acid –copper(II) and nickel(II) systems in ethanol-water mixture, *Ind. J. of chem.*, 42A, 2003, 1632-1635.
- Gandhi L. and Sekhon B.S., Complexes of silver picrate with Succinyl sulfathiazole and Sulfisoxazole, *International Journal of ChemTech Research*, Vol.2, No.1, 2010, 303-306.
- Erzalina Hernowo, Artik Elisa Angkawijaya, Ahmed E. Fazary, Suryadi Ismadji, and Yi-Hsu Ju\*, Complex Stability and Molecular Structure Studies of Divalent Metal Ion with L-Norleucine and Vitamin B3, *J. Chem. Eng. Data*, 56, 2011, 4549-4555.
- Pauwels T. F., Lippens W., Smet P. W., Herman G. G., Goeminne A. M., The dinuclear copper(II) complex of the hexaazamacrocyclic ligand 3,6,9,17,20,23-hexaazatriacyclo[23.3.1.1] triaconta-1(29), 11(30),12, 14,25,27-hexaene as a host for alanine, valine, leucine, norleucine, norvaline and serine anions, *Polyhedron*, 18(7),1999,1029-1037.
- Sanna D., Agoston C.G., Sovago I., Micera G., "Potentiometric and spectroscopic studies on the copper(II) complexes formed by oligopeptides containing histidine with a protection at the terminal amino group", *Polyhedron*, 20, 2001, 937-947.



8. Nelson G.A., Crawford M.B., Geddes B. J., Thiocarboxylates of nickel(II), *Inorg.Chem.*, 9,1970, 1123-1126.
9. Sayce I.G., computer calculation of equilibrium constants of species present in mixtures of metal ions and complexing agents, *Talanta*, 15, 1968, 1397-1411.
10. Sayce I.G., Sharma V. S, Computer calculation of equilibrium constants using programme SCOGS: a further modification, *Talanta*, 19(6), 1972, 831.
11. Thomas J. Difeo and Judith E. Shuster, Analytical profiles of drug substances and excipients - Vol. 23, *Brittain*, 1994,229.
12. Conner C.S., *Drug Intell. Clin. Pharm.*, 17, 1983, 898-899.
13. U.S.Pharmacopoeia, The National Formulary, Asian Edn, USP 24,NF 19, 2000, 867.
14. The Merck index, 12<sup>th</sup>Edn, White House Station, N.J., 1996, 49, 69.
15. Welling P.G. J. *Pharmacokinet*, *Biopharm*, 5, 1977, 291.
16. Williams D.A. and Lemke L.T., Foye's principles of medicinal chemistry, 5<sup>th</sup>Edn, 2005, 524- 525.
17. Maslowska J. and Chruscinski L, Potentiometric studies on mixed-ligand complexes of Cr(III) with dl-methionine, dl-ethionine and other amino acids, *Polyhedron*, 5(5), 1986, 1135-1139.
18. Artik Elisa Angkawijaya, Ahmed E. Fazary, Erzalina Hernowo, Mohamed Taha, and Yi-Hsu Ju, Iron(III), Chromium(III), and Copper(II) Complexes of L-Norvaline and Ferulic Acid, *J. Chem. Eng. Data*, 56, 2011, 532–540.
19. Joshi J. D., Vora J.J., Sharma Sangita, Gurjar J.G., Patel R.A., Patel D.R., Chaudhary R.H., Equilibrium studies on binary chelate formation of vitamin-U with some transition metal ions,*Int. J.Chem. Sci.*, 1(2), 2003, 155-158.
20. Ahmed EidFazary, Mohamed Taha, and Yi-Hsu Ju, Iron Complexation Studies of Gallic Acid, *J. Chem. Eng. Data*, 54, 2009, 35–42.
21. Sigel H., *Metal ions in biological systems*, Marcel Dekker, New York, 2, 1973, 63.
22. Wellman K.M., Mecca T.G., Mungall W., Hare C.R., The detection of apical interaction in copper (II) complexes of potential tridentate alpha-amino acids by optical rotatory dispersion., *J. Am. Chem. Soc.*, 90, 1968, 805-7.
23. LeussingD.L., Schiff base complexes: A numerical study of the nickel-pyruvateglycinate system using a high speed computer, *Talanta*, 11(2), 1964, 189-201.

\*\*\*\*\*

