

EVALUATION OF ANTI-PYRETIC POTENTIAL OF *FICUS CARICA* LEAVES

Patil Vikas V.*, Bhangale S.C., Patil V. R.

Department of Pharmaceutical Chemistry and Pharmacognosy
TVES's H'ble Loksevak Madhukarrao Chaudhari College of Pharmacy
Faizpur, Maharashtra

*Email: vikas312@rediff.com

ABSTRACT

A study was carried out to evaluate the anti-pyretic effect of an ethanol extract of leaves, of *Ficus carica* Linn. belonging to the family of *Moraceae*, at normal body temperature and yeast-induced pyrexia, in albino rats. A yeast suspension (10 ml/kg body wt.) increased rectal temperature 19 hours after the subcutaneous injection. The ethanol extract of *Ficus carica*, at doses of 100, 200 and 300 mg/kg body wt. p.o., showed significant dose-dependent reduction in normal body temperature and yeast-provoked elevated temperature. The effect extended up to five hours after drug administration. The anti-pyretic effect of the ethanol extract of *Ficus carica* was comparable to that of Paracetamol (150 mg/kg body wt., p.o.), a standard anti-pyretic agent.

Keywords: *Ficus carica*, Anti-pyretic effect, Ethanolic extract, Leaves, *Moraceae*, Yeast-induced pyrexia, Paracetamol.

INTRODUCTION

Ficus carica Linn. (Mar, Hindi & Guj: Anjir) Belongs to family *Moraceae*. A small or moderate sized deciduous tree, 15-30 ft high with broad ovate or nearly orbicular leaves, more or less deeply 3-5 lobed, rough above and pubescent below; fruits axillary, usually pear shaped, variable in size and colour.

The fig plant is considered to be a native of *carica* in asia and is grown in nearly all tropical and sub-tropical countries. It is now cultivated chiefly in the Mediterranean region, from Turkey in the east to Spain and Portugal in the west; it is also grown commercially in parts of U.S.A. and Chile and, to a small extent, in India, Arabia China and Japan.¹

Ficus carica Linn. (Syn: *Ficus sycomorus*; family: *Moraceae*) is commonly referred as "Fig". Its fruit, root and leaves are used in the native system of medicine in different disorders such as gastrointestinal (colic, indigestion, loss of appetite and diarrhea), respiratory (sore throats, coughs and bronchial problems), inflammatory and cardiovascular disorders^{2,3}. Fig has been traditionally used for its medicinal benefits as metabolic, cardiovascular, respiratory, antispasmodic and anti-inflammatory remedy^{4,5}.

Previous reports concerning the nutrient composition of dried figs have indicated that it has the best nutrient score among the dried fruit, being an important source of minerals and vitamins⁶.

The presence of Phytosterols (433 mg/100 g dry basis) has also been reported in fig fruit⁷. The fresh and dried figs also present relatively high amounts of crude fiber (5.5%, w/w) and polyphenols^{8,9}. Some recent works have reported that fig antioxidants can protect lipoproteins in plasma from oxidation and produce a significant increase in plasma antioxidant capacity for 4 h after consumption⁹. Also, showed that the higher the Polyphenols contents, especially Anthocyanins, in fig fruit, the higher was their antioxidant activity¹⁰.

MATERIALS AND METHODS

Plant material

Ficus carica Linn. leaves were collected from the Pal Satpuda ranges, in the Jalgaon district of Maharashtra, India, and identified by the Botanical Survey of India, Pune. The voucher specimen has been retained in our laboratory for future reference no. (BSI/WC/Tech/2008/355). The seeds were dried under controlled temperature, powdered and passed through a 40-mesh sieve.

Extraction Procedure

The powdered plant material was extracted using 90% ethanol, because ethanol extract is generally effective for the activity and also various chemical constituents may be present in the ethanol extract. The solvent was completely removed by vacuum distillation to yield a reddish-brown residue (yield 9.8%, w/w, with respect to dry starting material). This ethanolic extract of *Ficus carica* was examined chemically and was observed to contain flavonoids, steroids, triterpenoids, alkaloids and tannins. These constituents were confirmed using thin-layer chromatography (TLC). A weighed amount of the dried ethanol extract of the seeds of *Ficus carica* was suspended in 2% aqueous Tween 80 solution and used for the present study.

Animals Used

Adult albino rats (Wistar strain) of either sex, weighing 180-200 g each, were used. The animals were maintained under suitable nutritional and environmental conditions throughout the experiment.

Study on Normal Body Temperature

Rats of either sex were divided into four groups of six each. The body temperature of each rat was measured rectally at predetermined intervals before and five hours after administration of either 2% aqueous Tween 80 solution (control) or ethanol extract of *Ficus carica* at oral doses of 100, 200 and 300 mg/kg body wt.

Induction of Yeast-Induced Pyrexia

The rats were divided into five groups of six each. The normal body temperature of each rat was measured rectally at predetermined intervals and recorded¹¹. The rats were trained to remain quiet in a restraint cage. A thermistor probe was inserted 3–4 cm deep into the rectum and fastened to the tail by adhesive tape. Temperature was measured on a digital thermometer. After measuring the basal rectal temperature, the animals were injected subcutaneously with 10 ml/kg body wt. of 15% w/v yeast, suspended in 0.5% w/v methylcellulose solution. The rats were then returned to their housing cages. Nineteen hours after the yeast injection, the animals were again restrained in individual cages for rectal temperature recording, as described previously.

Drug Administration

Nineteen hours after yeast injection, the ethanol extract of *Ficus carica* was administered orally at doses of 100, 200 and 300 mg/kg body wt. to three groups of animals, respectively. A similar volume (5 ml/kg body wt.) of 2% aqueous Tween 80 solution was administered orally to the control group. The fifth group of animals received the standard drug, Paracetamol (150 mg/kg body wt.), orally. The rats were restrained for rectal temperature recording at the nineteenth hour, immediately before ethanol extract of *Ficus carica* or saline or Paracetamol administration, and again at one hour intervals up to the twenty-third hour after yeast injection.

Statistical Analysis

The data were analyzed for significance using the unpaired two-tailed Student's t-test¹².

RESULTS

The effect of the ethanol extract of *Ficus carica* on the normal body temperature in rats is shown in [Table I]. It was found that ethanol extract of *Ficus carica* at doses of 100 mg/kg body wt. caused significant lowering of body temperature up to the fourth hour following its administration. This effect was maximum at doses of 200 and 300 mg/kg body wt., in a dose-dependent manner, and caused significant lowering of body temperature up to the fifth hour after its administration. The subcutaneous injection of yeast suspension markedly elevated the rectal temperature at the nineteenth hour after administration. Treatment with ethanol extract of *Ficus carica* at doses of 100, 200 and 300 mg/kg body wt. decreased the rectal temperature of the rats in a dose dependent manner. The antipyretic effect started as early as the first hour after administration, and the effect was maintained for four hours after its administration. The standard drug Paracetamol at the 150 mg/kg body wt. dose significantly reduced the yeast-provoked elevation of body temperature. The results obtained for both the standard drug-treated and ethanol extract of *Ficus carica* treated rats were compared with the control (2% aqueous Tween 80 solution) group and a significant reduction in the yeast-elevated rectal temperature was observed [Table II].

Table I: Effect of Ethanolic Extract of *Ficus carica* (EEFC) on normal body temperature

Treatment	Rectal temperature (°C) before and after treatment					
	0h	1h	2h	3h	4h	5h
Control 5ml/kg body wt.	27.4±0.2	27.21±0.1	27.38±0.31	27.12±0.2	27.12±0.1	27±0.2
EEFC 100mg/kg body wt.	27.2±0.1	26.7±0.3 ^c	26.6±0.2 ^b	26.5±0.3 ^b	26.7±0.2 ^b	26.8±0.3
EEFC 200mg/kg body wt.	27.3±0.2	26.2±0.1 ^a	26.2±0.2 ^a	26.3±0.2 ^b	26.7±0.2 ^b	26.7±0.2 ^b
EEFC 300mg/kg body wt.	27.1±0.2	25.8±0.2 ^a	25.8±0.2 ^a	25.8±0.1 ^a	25.9±0.2 ^a	25.0±0.2 ^b

Each value represents mean ± SEM (n=6)

Control = 2% aqueous Tween 80 solution

^ap<0.001, ^bp<0.01, ^cp<0.05, as compared to control values for the corresponding hour.

EEFC - Ethanolic Extract of the leaves of *Ficus carica* Linn.

Table II: Effect of Ethanolic Extract of *Ficus carica* (EEFC) on yeast-induced pyrexia in rats

Treatment	Rectal temperature (°C) before and after treatment					
	0h	19h	20h	21h	22h	23h
Control 5ml/kg body wt.	27.59±0.02	29.7±0.02	29.49±0.03	29.3±0.07	29.0±0.03	29.05±0.03
Paracetamol 150mg/kg body wt.	27.9±0.01	29.7±0.03	28.1±0.01 ^a	28.0±0.01 ^a	27.9±0.05 ^a	27.8±0.03 ^a
EEFC 100mg/kg body wt.	27.4±0.04	29.8±0.01	29.1±0.01 ^b	28.49±0.02 ^b	28.3±0.05 ^b	27.7±0.08 ^a
EEFC 200mg/kg body wt.	27.4±0.01	29.8±0.01	28.8±0.05 ^a	28.3±0.05 ^a	27.9±0.08 ^a	27.7±0.06 ^a
EEFC 300mg/kg body wt.	27.5±0.07	29.8±0.07 ^a	28.5±0.01 ^a	27.8±0.03 ^a	27.4±0.01 ^a	27.49±0.05 ^a

Each value represents mean ± SEM (n=6)

Control = 2% aqueous Tween 80 solution

^ap<0.001, ^bp<0.01, as compared to control values for the corresponding hour.

EEFC - Ethanolic Extract of the leaves of *Ficus carica* Linn.

DISCUSSION

Fever may be a result of infection or one of the sequels of tissue damage, inflammation, graft rejection, or other disease states. Antipyretics are drugs which reduce the elevated body temperature. Regulation of body temperature requires a delicate balance between the population and loss of heat, and the hypothalamus regulates the set point at which body temperature is maintained. In fever, this set point is elevated, and drugs like Paracetamol do not influence body temperature when it is elevated by factors such as exercise or increase in ambient temperature¹³.

In the present study the effects of various concentrations of plant extracts and the effects of extracts with that of Paracetamol at different times are compared. The present result shows that the ethanolic extract of the leaves of *Ficus carica* possesses a significant antipyretic effect in the yeast-provoked elevation of body temperature in rats, and its effect is comparable to that of Paracetamol (standard drug). Furthermore, the ethanol extract of *Ficus carica* also significantly reduces the normal body temperature, and this is to be studied further for the exact mechanism of action.

REFERENCES:

1. The Wealth of India Volume (F-G) In : A Dictionary of Indian Raw Materials and Industrial Products Vol. 4 New Delhi: council of Scientific and Industrial Research;1999;27.
2. Burkill, I.H., A Dictionary of the Economic Products of Malay Peninsular. Ministry of Agriculture, Malaysia, 1935;1005–1006.
3. Ponelope O., 100 Great Natural Remedies Kyle Cathic Limited, NewYork, USA, 1997;98-99.
4. Duke, J.A., Bogenschutz-Godwin, M.J., Du Celliar, J., Duke, P.K., Hand Book of Medicinal Herbs, second ed. CRC Press, Boca Raton, USA, 2002; 314–315.
5. Werbach M., Healing with Food. Harper Colines, NewYork USA,1993;443-444.
6. US Department of Agriculture, Agricultural Research Service, USDA Nutrient Database for Standard Reference, Release 15, (2002) Nutrient Data Laboratory Home Page, www.nal.usda.gov/fnic/foodcompS.
7. Jeong, W.S., Lachance, P.A., Phytosterols and fatty acids in fig (*Ficus carica*, var. Mission) fruit and tree components. Food Chemistry and Toxicology 2001;66: 278–281.
8. Vinson, J.A., The functional food properties of figs. Cereal Foods World 4, 1999; 82–87.
9. Vinson, J.A., Zubik, L., Bose, P., Samman, N., Proch, J., Dried fruits: excellent in vitro and in vivo antioxidants. Journal of the American College of Nutrition 2005;4, 44–50.
10. Solomon, A., Golubowicz, S., Yablowicz, Z., Grossman, S., Bergman, M., Gottlieb, H.E., Altman, A., Kerem, Z., Flaishman, M.A., Antioxidant activities and anthocyanin content of fresh fruits of common fig (*Ficus carica* L.). Journal of Agricultural and Food Chemistry 2006; 54:7717–7723.
11. Vogel HG, editor. Drug Discovery and Evaluation Pharmacological Assays 2nd ed. New York: Springer-Verlager Berlin Heidelberg; 2002;696-7,692.
12. Woodson RF. Statistical methods for the analysis of biochemical data, in probability and mathematical statistics. Wiley, New York, Chichester: 1987; 315-6.
13. Goodman, Gilman. The Pharmacological Basis of Therapeutics. 9th ed. New York: McGraw-Hill, Professions division; 1996; 959-75.
